

Contents

1	Introduction	2
2	Coherent States and the Coherent State Path Integral	6
2.1	Canonical coherent states	6
2.2	General coherent states	9
2.3	Group defined coherent states	10
2.4	The standard coherent state path integral	17
2.5	Mathematical aspects of path integration	20
2.6	Mathematical tools	22
2.7	The coherent state path integral with Wiener measure	29
3	Weak Coherent States and the Weak Coherent State Path Integral	37
3.1	Weak coherent states	38
3.2	The affine weak coherent state path integral	39
3.2.1	Spectral approach	40
3.2.2	Introducing dynamics	61
3.2.3	Regularizing approach	64
3.2.4	Introducing dynamics	77
4	Final Remarks	80
A	Appendix	82
A.1	The affine group	82
A.2	Parameters in DKP and the present work	83
A.3	Spectrum of A	85
A.4	Alternative computation of the denominator of Eq. (75)	95
B	Acknowledgments	99

1 Introduction

In standard quantum mechanics, in the Schrödinger or Heisenberg formulation, phase space and its methods - intensely studied in classical mechanics - play no role. In the quantization procedure the classical variables q (space) and p (momentum) are promoted to operators Q and P satisfying the commutation relation $[Q, P] = i$, where throughout the article $\hbar = 1$. These act on a Hilbert space of state vectors in the well-known way. Formally there is a strong resemblance between classical dynamics formulated in terms of Poisson brackets and quantum dynamics formulated with commutators (Heisenberg picture). But the classical Heisenberg equations of motion are covariant under canonical transformations, a property related to the symplectic structure of the underlying phase space manifold, i.e. a geometric structure. On the other hand the canonical quantization procedure works only in one special coordinate system, namely Cartesian coordinates, as was pointed out by Dirac[1]¹. This certainly is a dilemma, since quantization should not depend on something so arbitrary as the choice of coordinates. And moreover Cartesian coordinates exist only in flat space.

This was fully recognized and many efforts were made to reformulate quantum mechanics in phase space. This led e.g. to the so-called deformation quantization and to geometric quantization schemes, i.e. quantizations which respect the free choice of coordinates and allow canonical transformations. A short introduction to the general ideas of these quantizations will subsequently be given since some of their ideas are borrowed in the present work.

Deformation Quantization

The oldest formulation of quantum mechanics in phase space is due to Wigner [3][4], Moyal [5] and others [6][7][8][9][10][11].

Wigner's idea was to introduce a quasi-probability function² on phase space (here 2-dimensional for sake of simplicity)

$$F(p, q) = \frac{1}{2\pi} \int \psi^*(q - x/2) e^{-ixp} \psi(q + x/2) dx$$

which has the properties $\int F(p, q) dp = |\psi(q)|^2$ and $\int F(p, q) dq = |\tilde{\psi}(p)|^2$

¹For an example that one can obtain wrong results in other coordinate systems see e.g. [2]

²Here "quasi-probability" refers to the fact that these functions are not positive everywhere for general ψ .

giving the right quantum mechanical expectations for the state $|\psi\rangle$ in the q - and p -representation respectively.

Moyal [5] showed that when quantum mechanics is expressed in phase space in this way quantum dynamics (and therefore the analog of the classical Poisson bracket) is governed by the Moyal sine-bracket, which is the quantum deformed version of the Poisson bracket and the Moyal equations of motion - analog to the Liouville equations and with $H(p, q)$ the classical Hamiltonian - become:

$$\partial_t F(p, q, t) = 2 \sin\{\frac{1}{2}[\partial_{p_F} \partial_{q_H} - \partial_{p_H} \partial_{q_F}]\} H(p, q) F(p, q, t)$$

In the same paper Moyal also showed the equivalence of this statistical approach to quantum mechanics to the usual Schrödinger formulation (the Wigner function is a special representation of the density matrix in the so-called Weyl-correspondence, see e.g. [11]). The more modern notation uses the $\star$ -product, defined as $\star := \exp\{\frac{i}{2}(\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x)\}$, where the arrows indicate the side this operator acts on, and the Moyal equations become $i\partial_t F = H \star F - F \star H$. Since the Schrödinger and this phase space formulation are equivalent it is not surprising that things like Heisenberg uncertainty relation etc. emerge naturally ([11]), but on the other hand this means that it is not surprising either that canonical transformations do not in general leave the $\star$ -product invariant³.

An interesting point is the connection of the Wigner functions (or more precisely of their Fourier transform) to coherent states. This will be discussed later after having introduced the concept of coherent states.

Geometric Quantization

Geometric quantization tries - as the name already implies - to formulate the quantization procedure in such a way that it is fully geometric in character. The phase space M is taken as the starting point and functions $\psi(p, q)$, $(p, q) \in M$, are introduced which are supposed to be square integrable. The basic quantum kinematical operators Q and P are then represented with linear differential operators according to some rule, e.g. $Q = i\partial_p + q/2$, $P = -i\partial_q + p/2$, the only requirement being that they fulfill the Heisenberg commutation relation $[Q, P] = i$. But there is a problem since these representations are reducible⁴ and this very fact - although non-critical from a

³Only linear canonical transformations do, [11]

⁴For the example given above the operators $\tilde{Q} = i\partial_p - q/2$ and $\tilde{P} = -i\partial_q - p/2$ commute with Q and P and are certainly not multiples of the identity.

mathematical point of view - simply leads to inconsistencies with physics. So the whole process up to this point was eventually called pre-quantization.

The idea to remedy the situation is “polarization” and initially it came from the fact that the Schrödinger wave functions depend on q only (and the operators $P = -i\partial_q$ and $Q = q$ are irreducibly represented). A real polarization $\partial_p\psi(p, q) = 0$ would lead to functions which depend on q only but not to square integrable functions of phase space. So complex linear polarizations, say $[\partial_q - i(\partial_p + p)]\psi(p, q) = 0$ just to give an example, were imposed to reduce the space of admissible functions since $L^2(M)$ was too big to lead to irreducible representations of the basic kinematical operators Q and P . Finally, by one way or another, schemes to deal with kinematics could be established which were also consistent with the kinematical versions of (standard) quantum mechanics⁵. The breakdown of the theory came with dynamics. The idea was to copy the classical Poisson bracket and establish an operator formalism which respects the same relations (thus conserving the geometric character of the classical world). Suffice it to say that the rules derived in this way simply led to wrong answers except for linear Hamiltonians (and by chance for the harmonic oscillator) and the whole program was largely abandoned.

The important connection for the present work is the idea of polarization. As will be shown some coherent states already have a linear complex polarization condition built into them by their very definition (and they automatically form a subset of $L^2(M)$ which leads to irreducible representations of the kinematical operators).

Coherent States and their Path Integrals

Coherence is a very general concept and especially important in the field of optics where it was studied for more than three centuries. It is thus not surprising that the first coherent states, nowadays called canonical coherent states, arose in quantum optics [13]. The name “coherent states” was chosen as an analog to the classical coherent ensemble, i.e. an ensemble for which $\langle V^*(t_1)\dots V^*(t_r)V(t_{r+1})\dots V(t_{2r}) \rangle = Y^*(t_1)\dots Y^*(t_r)Y(t_{r+1})\dots Y(t_{2r})$, where $\langle \dots \rangle$ denotes the ensemble average and Y is a deterministic function. A (canonical) coherent state $|\psi\rangle$ is a state for which similarly $\langle \psi|a^\dagger(t_1)\dots a^\dagger(t_r)a(t_{r+1})\dots a(t_{2r})|\psi\rangle = z^*(t_1)\dots z^*(t_r)z(t_{r+1})\dots z(t_{2r})$, where a and $a^\dagger$ are the harmonic oscillator creation and annihilation operators and z is a complex valued function. Soon the concept of coherent states gradually broadened.

⁵For further reading see e.g. [12]

Since they will be extensively discussed in the following chapter it may suffice at this point to say that, generally speaking, they are by their very definition related to a certain underlying phase space. Hence they are a third way to describe quantum mechanics in such a way that phase space methods can be used (e.g. canonical transformations of the underlying phase space do not change the set of coherent states associated with it). A quantization by path integrals based on coherent states thus promises to be of a geometric nature once all difficulties with well-definedness are overcome. Indeed, such a path integral involving a metric regularization that leads to a so-called Wiener measure has been developed by I. Daubechies and J.R. Klauder and T. Paul, and this result is the basis to understand the present work which studies the existence of a similar path integral for even more general states. These states are called weak coherent states and together with the coherent states they form the class of Klauder states.

2 Coherent States and the Coherent State Path Integral

Initially, the term “coherent states” was used in a quite narrow sense, namely what are called “canonical” coherent states today. Gradually, the concept was extended, and “generalized” coherent states or “Overcomplete Families of States” were used in many applications. Klauder was the first to lay open the fundamental properties they share, and they are all subsumed in the sense of his very general definition as coherent states nowadays.

For a better understanding and to get a feeling for coherent states this mathematical framework is not taken as a starting point for section (2.1), but rather the original canonical coherent states. Many interesting properties of coherent states can already be learned in the discussion of this example. This also makes it easier to understand the general concept in the following section (2.2) which otherwise might seem quite technical. Next, section (2.3) specializes again to so-called group-defined coherent states which will be of most interest in the subsequent work. The ultimate goal of the whole chapter is to establish a well-defined coherent state path integral which will provide a quantization scheme fully geometric in nature. To this end the standard coherent state path integral is presented first in section (2.4) and its shortcomings are pointed out in section (2.5), followed by section (2.6) containing mathematical tools which prepare the ground for the understanding of the final coherent state path integral with Wiener measure [section (2.7)].

2.1 Canonical coherent states

Still the best known and most widely used coherent states are the canonical coherent states.

Let $a, a^\dagger$ be operators obeying the Heisenberg algebra, i.e. $[a, a^\dagger] = 1$, and let $|0\rangle$ be the unique normalized state annihilated by a , i.e. $a|0\rangle = 0$. Denote the eigenstates of the number operator $N = a^\dagger a$ by $|n\rangle := \frac{1}{\sqrt{n!}}(a^\dagger)^n|0\rangle$. Then the coherent states are defined as [14]

$$\begin{aligned} |z\rangle &:= e^{za^\dagger - z^*a}|0\rangle = e^{-\frac{1}{2}|z|^2} e^{za^\dagger} e^{-z^*a}|0\rangle \\ &= e^{-\frac{1}{2}|z|^2} e^{za^\dagger}|0\rangle = e^{-\frac{1}{2}|z|^2} \sum_{n=0}^{\infty} \frac{1}{\sqrt{n!}} z^n |n\rangle \end{aligned} \quad (1)$$

The relation $ae^{za^\dagger} = e^{za^\dagger}e^{-za^\dagger}ae^{za^\dagger} = e^{za^\dagger}(a+z)$ immediately follows and

applying this to the second but last equality in (1) it is easily seen that $a|z\rangle = z|z\rangle$, i.e. the canonical coherent states are the generalized eigenstates of the annihilation (or lowering) operator a . From Eq. (1) the overlap is easily seen to be

$$\langle z_2|z_1\rangle = e^{-\frac{1}{2}|z_1|^2 - \frac{1}{2}|z_2|^2 + z_2^* z_1} \quad (2)$$

which is a jointly continuous, nowhere vanishing function of z_1 and z_2 .

By the foregoing $\langle z|a|z\rangle = z\langle z|z\rangle = z$, so the label z is the mean of the operator a in the coherent state $|z\rangle$.

Consequently each state $|z\rangle$ satisfies $\langle z|a^\dagger a|z\rangle = |z|^2 = |\langle z|a|z\rangle|^2$ and since the uncertainty relation for a and $a^\dagger$ simply is $\langle a^\dagger a\rangle \geq \langle a^\dagger\rangle\langle a\rangle$ every single canonical coherent state is a minimum uncertainty state.

The vector-valued function $|z\rangle$ is continuous in z [i.e., if $z \rightarrow z'$ then $\| |z\rangle - |z'\rangle \| \rightarrow 0$] and the complex-valued function $\psi(z) = \langle z|\psi\rangle$ is bounded and continuous in z for all $|\psi\rangle \in$ Hilbert space. Further $\psi(z) = 0 \Leftrightarrow |\psi\rangle = 0$ holds establishing a one-to-one relation between functions $\psi(z)$ and vectors $|\psi\rangle$, and thus $\psi(z)$ can be taken as a representative of the vector $|\psi\rangle$.

With the help of the last expression in Eq. (1) and the use of polar coordinates at an intermediate step it can be seen that the canonical coherent states resolve unity in the form

$$\begin{aligned} \pi^{-1} \int |z\rangle\langle z| d^2 z &:= \pi^{-1} \sum_{n,m} \frac{1}{\sqrt{n!m!}} \int e^{-|z|^2} z^{*n} z^m |m\rangle\langle n| d[\Re(z)] d[\Im(z)] \\ &= \sum_n \frac{1}{n!} \int e^{-|z|^2} |z|^{2n} |n\rangle\langle n| d[|z|^2] = \sum_n |n\rangle\langle n| = \mathbb{1} \end{aligned} \quad (3)$$

Note that a resolution of unity guarantees the existence of a coherent state representation of Hilbert space (it is a stronger property than $\psi(z) = 0 \Leftrightarrow |\psi\rangle = 0$ which was already sufficient for this purpose).

Since the canonical coherent states depend only on z but not on z^* the functions $\psi(z) := \langle z|\psi\rangle = e^{-|z|^2/2} f(z^*)$ in the coherent state representation are entire functions $f(z^*)$ apart from an overall factor. This shows that the canonical coherent states are overcomplete (especially they can not all be orthogonal) since by analyticity the vanishing of $\psi(z)$ on a characteristic set (e.g. any curve of nonzero length in the complex plane) already implies that $\psi(z) \equiv 0$.

By analyticity one has $\partial_z(e^{|z|^2/2}\langle z|\psi\rangle) = e^{|z|^2/2}(\partial_z + z^*/2)\langle z|\psi\rangle = 0$ and thus $(\partial_z + z^*/2)\langle z|\psi\rangle = 0$ which is a complex polarization condition. Note

that this is really a consequence of $a|0\rangle = 0$ since $(\partial_z + z^*/2)\langle z|\psi\rangle = \langle 0|(\partial_z + z^*/2)e^{-\frac{1}{2}|z|^2}e^{za^\dagger}|\psi\rangle = \langle 0|-a^\dagger e^{-\frac{1}{2}|z|^2}e^{za^\dagger}|\psi\rangle = 0$.

A further consequence is that an operator B is already determined by its diagonal coherent state matrix elements $B(z) := \langle z|B|z\rangle = \exp\{-|z|^2\} \sum_{n,m} 1/\sqrt{n!m!} \cdot \langle n|B|m\rangle z^{*n} z^m$ since every monomial $z^{*n} z^m$ can be isolated by its dependence on θ and r (with $z := re^{i\theta}$) and analytically extended to $z^{*n} z'^m$ resulting in the general matrix element $\langle z|B|z'\rangle$. The diagonal matrix element $B(z)$ as a function of z is called the upper symbol or in this case of canonical coherent states the normal ordered symbol.

Additionally there are two other operator representations. The first is called diagonal representation and is given by $B = \pi^{-1} \int b(z)|z\rangle\langle z|d^2z$. The function $b(z)$ is called the lower symbol or in the case of canonical coherent states the anti-normal ordered symbol⁶.

The second is a differential operator representation. The creation and annihilation operators e.g. are represented by $(a^\dagger \leftrightarrow z^*)$ and $(a \leftrightarrow z/2 + \partial_{z^*})$ since $\langle z|a^\dagger|\psi\rangle = z^*\langle z|\psi\rangle$ and $\langle z|a|\psi\rangle = (z/2 + \partial_{z^*})\langle z|\psi\rangle$. A general operator $B(a^\dagger, a)$ is consequently represented by $B(z^*, z/2 + \partial_{z^*})$.

To show just a glimpse of the power of coherent states consider the so-called Segal-Bargmann (or holomorphic) representation of Hilbert space. Let $z^* = \zeta$ and let the normalization factor be absorbed into the measure $d\mu(\zeta) = \pi^{-1} \exp\{-|\zeta|^2\}d^2\zeta$. Then $\psi(\zeta) := \langle \zeta|\psi\rangle$ are entire (holomorphic) functions with inner product $\langle \psi|\psi\rangle = \int |\psi(\zeta)|^2 d\mu(\zeta)$ on the Hilbert space. In this form the operators $a^\dagger$ and a are represented by ζ and ∂_ζ respectively. The eigenvalue equation for the number operator $N = a^\dagger a$ becomes $\zeta \partial_\zeta \psi(\zeta) = \lambda \psi(\zeta)$ with solutions $\psi(\zeta) = c\zeta^\lambda$. But by analyticity $\lambda \in \mathbb{N}_0$ and therefore the spectrum of the number operator is derived in just these few lines!

Another example of the outstanding properties of the canonical coherent states is the stability under time evolution in a harmonic potential: $\exp\{-it\omega a^\dagger a\}|z\rangle = \exp\{-|z|^2/2 \sum_n \frac{1}{\sqrt{n!}} z^n \exp\{-in\omega t\}|n\rangle = |\exp\{-i\omega t\}z\rangle$, and since all canonical coherent states are minimum uncertainty states this property is preserved for all times.

For a wide range of applications of the canonical coherent states and the general coherent states which will be defined in the following section, see

⁶Many authors use the terminology exactly the other way round and call the upper symbol the lower one and vice versa. The reason is that the upper symbol (their lower symbol) is involved in a lower bound in certain inequalities, the Berezin-Lieb inequalities, and the lower symbol (their upper symbol) is involved in an upper bound in the same inequalities.

e.g. [14].

2.2 General coherent states

The definition of coherent states used here follows Klauder [14]. Let $\mathfrak{L}$ be a topological space called the “label space” and for $l \in \mathfrak{L}$ denote by $|l\rangle$ a vector in a Hilbert space $\mathfrak{H}$. For $|l\rangle$ to be called a coherent state two properties are required:

1) Continuity: The states $|l\rangle$ are a strongly continuous vector-valued function of the label l .

2) Resolution of unity: There is a positive measure δl on the label space $\mathfrak{L}$ such that the identity operator $\mathbb{1}$ on $\mathfrak{H}$ can upon integration over $\mathfrak{L}$ be represented as

$$\mathbb{1} = \int |l\rangle\langle l| \delta l \quad (4)$$

Condition 1) says: If $l' \rightarrow l$ in $\mathfrak{L}$ then $\| |l'\rangle - |l\rangle \| \rightarrow 0$ in $\mathfrak{H}$. Hence, the set $\{|l\rangle : l \in \mathfrak{L}\}$ is a continuous submanifold of $\mathfrak{H}$ if $\mathfrak{L}$ is connected.

Condition 2) is to be understood in the sense of weak convergence, i.e. arbitrary matrix elements converge in $\mathbb{C}$. Condition 2) implies completeness of the set $\{|l\rangle : l \in \mathfrak{L}\}$. Without loss of generality (w.l.o.g.) all $|l\rangle \in \mathfrak{H}$ can be assumed to be normalized to 1 since the measure δl on $\mathfrak{L}$ can be rescaled at every point to allow for this choice. Unless stated otherwise this convention will be adopted throughout the article.

To get a better feeling of what coherent states “are” it is certainly instructive to see what they are not. They can not be a set of discrete orthonormal vectors $\{|n\rangle : n \in \mathbb{N}\}$ which is not continuous in the labels and they can not be a set of δ -orthonormalized continuum vectors $\{|x\rangle : x \in \mathbb{R}\}$ which are continuous in the labels but do not form a continuous set (apart from the fact that they are not vectors in Hilbert space, but so-called generalized vectors). In other terms, coherent states can not be the eigenvectors (in the usual and generalized sense) of any self adjoint operator.

Also there is a difference in the resolution of unity afforded by the coherent states to the usual one furnished by generalized eigenvectors of self-adjoint operators (and there should be since the latter always exists. So it would not make sense to include the resolution of unity in the defining properties for coherent states if there wasn’t more to it). The one-dimensional

projection operators $|l\rangle\langle l|$ in Eq. (4) are not in general mutually orthogonal. The resolution of unity gives rise to a functional representation

$$\langle l|\phi\rangle = \int \langle l|l'\rangle\langle l'|\phi\rangle\delta l' \quad (5)$$

For δ -orthonormalized continuum vectors an equation of this form would merely be an identity, but because of the continuity requirement for coherent states the kernel $\langle l|l'\rangle$ is a jointly continuous function of l and l' , nonzero for $l = l'$ (for the normalized coherent states the value at this point is 1) and therefore nonzero in an open neighborhood due to continuity. This implies severe restrictions on the admissible functions. One speaks of a reproducing kernel and the subspace spanned by the coherent states is itself a so-called reproducing kernel Hilbert space (see e.g. [23]).

The coherent states are linearly dependent as is evident from $|l'\rangle = \int |l\rangle\langle l|l'\rangle\delta l$. This is why they are often called an “overcomplete” family of states.

2.3 Group defined coherent states

In this work group related coherent states will play an important role. These are defined in the following way [14][15]:

Let G be a Lie group and $U(g)$, $g \in G$, a strongly continuous irreducible unitary representation on a Hilbert space $\mathfrak{H}$. Let $|\psi_0\rangle$ be an arbitrary but fixed normalized vector in $\mathfrak{H}$ called the “fiducial” vector, then the action of U on $|\psi_0\rangle$ creates the coherent states of the group G :

$$|g\rangle := U(g)|\psi_0\rangle \quad (6)$$

where g runs over the whole group.

Assume for a moment that the group has n infinitesimal self-adjoint generators L_i , $i \in \{1, \dots, n\}$, and let $l \in \mathcal{L}$ be an element of an n -dimensional “label” space (a topological space) such that the group is represented in group coordinates of the second kind as $U(l) = e^{il_1 L_1} \dots e^{il_n L_n}$ which is supposed to be irreducible. Then we call the label space the phase space associated with the group. Under canonical transformations of the phase space variables such that $l_1 = l_1(\bar{l}_1, \dots, \bar{l}_n)$, $\dots$, $l_n = l_n(\bar{l}_1, \dots, \bar{l}_n)$ and with an arbitrary phase $\bar{F}_0(\bar{l}_1, \dots, \bar{l}_n)$ it follows that

$$|l_1 \dots l_n\rangle \rightarrow \overline{|\bar{l}_1 \dots \bar{l}_n\rangle} := e^{-i\bar{F}_0(\bar{l}_1, \dots, \bar{l}_n)} |l_1(\bar{l}_1, \dots, \bar{l}_n) \dots l_n(\bar{l}_1, \dots, \bar{l}_n)\rangle \quad (7)$$

Apart from a phase factor the set of coherent state vectors has remained the same, the only difference being the way they are labeled. The “new” coherent states are still generated by the action of the group G on the fiducial vector $|\psi_0\rangle$ but the group is generally no longer represented in canonical group coordinates [28].

The overlap of two states $|g\rangle$ and $|g'\rangle$, $\langle g|g'\rangle = \langle\psi_0|U(g)^\dagger U(g')|\psi_0\rangle = \langle\psi_0|U(g^{-1} * g')|\psi_0\rangle$, is bounded (by unity) and jointly continuous in the group parameters.

Completeness of these states follows immediately from the irreducibility of U [15] but to qualify as coherent states they must admit a resolution of unity in the form $\mathbb{1} = \int |l\rangle\langle l| \delta l$. Klauder has shown [16][17] that the integration measure can be taken as the left-invariant group measure dg ⁷. If the representation U of G is square integrable, i.e. $\int |\langle\psi_0|U(g)|\psi_0\rangle|^2 dg < \infty$ ⁸, then - with the normalization $\int |\langle\psi_0|U(g)|\psi_0\rangle|^2 dg = 1$ - the invariance of the measure along with Schur’s Lemma provides a resolution of unity

$$\mathbb{1} = \int_G |g\rangle\langle g| dg \quad (8)$$

Next consider a general operator $\mathcal{H}$ on the Hilbert space $\mathfrak{H}$. Associated with it are the upper symbol

$$H(g) = \langle g|\mathcal{H}|g\rangle \quad (9)$$

which is the diagonal coherent state matrix element for given g , and (if it exists) the lower symbol implicitly defined as the diagonal representative of $\mathcal{H}$

$$\mathcal{H} = \int h(g) |g\rangle\langle g| dg \quad (10)$$

It was shown by Lieb ([14], page 35) that the set of operators admitting a diagonal representation, i.e. for which the lower symbol exists, is identical to the set of operators uniquely specified by the diagonal coherent state matrix elements alone, i.e. by the upper symbol.

If $\mathcal{H}$ is interpreted as the quantum Hamiltonian then each of the symbols is interpreted in various physical applications as the associated classical

⁷For compact groups left- and right-invariant measure always coincide. More generally left- and right-invariant group measure coincide if the structure constants of the group algebra satisfy $C_{ab}^b = 0$.

⁸This bound is automatic if the group volume is finite which is always the case for compact groups, but it can impose restrictions for non-compact groups.

Hamiltonian. For the upper symbol this association is known as the “weak correspondence principle” [17][22].

The symbols associated with an operator will be very important in the coherent state path integral formulation.

In chapter 3 this work will deal almost entirely with affine coherent states which are special coherent states generated by the affine group, often called the $ax+b$ -group, which is a subgroup of $SU(1,1)$. Nevertheless for illustration and for the sake of comparison it is useful to start with the most common coherent states, the canonical coherent states, which are the coherent states of the Heisenberg-Weyl group. After that example a short overview of the spin coherent states, the coherent states of the $SU(2)$ group, will follow before the affine coherent states are discussed.

Canonical coherent states - phase space formulation:

The Heisenberg-Weyl group is generated by the self-adjoint canonical operators P and Q satisfying the Heisenberg commutation relation $[Q, P] = i$ and has exactly one irreducible unitary representation up to unitary equivalence which is e.g. given in canonical group coordinates of the second kind by

$$U(p, q) = e^{-iqP} e^{ipQ} \quad (11)$$

where $(p, q) \in \mathbb{R} \times \mathbb{R}$.

The canonical coherent states are generated by the action of this group on the harmonic oscillator ground state $|0; \omega\rangle$ which satisfies $(\omega^{1/2}Q + i\omega^{-1/2}P)|0; \omega\rangle = 0$ ($\omega \in \mathbb{R}$ is the oscillator frequency). The ground state will often be written as $|0\rangle$ with the ω -dependence left implicit. As stated previously, the ground state is a minimum uncertainty state. The definition of the coherent states is

$$|pq\rangle := U(p, q)|0\rangle \quad (12)$$

The canonical coherent states introduced in the beginning of this chapter in their complex characterization are the same states apart from an unimportant phase factor. To see this rewrite the annihilation operator as $a = 2^{-1/2}(\omega^{1/2}Q + i\omega^{-1/2}P)$, the creation operator as $a^\dagger = 2^{-1/2}(\omega^{1/2}Q - i\omega^{-1/2}P)$ and the complex label z as $z := 2^{-1/2}(\omega^{1/2}q + i\omega^{-1/2}p)$. Then $U(z) := e^{za^\dagger - z^*a} = e^{i(pQ - qP)} =: \tilde{U}(p, q)$ and the group representation is given in canonical group coordinates of the first kind.

The overlap of two states defined by Eq. (12) is

$$\langle pq|p'q'\rangle = e^{i/2(p+p')(q-q')} e^{-1/4[\omega^{-1}(p-p')^2 + \omega(q-q')^2]} \quad (13)$$

whereas the phase would be different if the group had been defined with group coordinates of the first kind, namely $\langle pq|p'q'\rangle = \exp\{\frac{i}{2}(p'q - q'p)\} \exp\{-\frac{1}{4}[\omega^{-1}(p-p')^2 + \omega(q-q')^2]\}$. The Weyl operator U serves as a translation operator for P and Q

$$\begin{aligned} U^\dagger(p, q)QU(p, q) &= Q + q \\ U^\dagger(p, q)PU(p, q) &= P + p \end{aligned} \quad (14)$$

showing that the coherent states $|pq\rangle$ are labeled by the means of p and q since $\langle 0|Q|0\rangle = \langle 0|P|0\rangle = 0$ and so $\langle pq|Q|pq\rangle = q$ and $\langle pq|P|pq\rangle = p$.

The resolution of unity reads

$$\mathbb{I} = (2\pi)^{-1} \int |pq\rangle\langle pq| dp dq \quad (15)$$

Since the group measure is both left- and right-invariant, this implies that all fiducial vectors lead to square-integrable representations if just one vector does [14]. And clearly there is one such vector, namely the harmonic oscillator ground state $|0\rangle$.

General bounded operators, or operators polynomial in P and Q , are uniquely determined by their diagonal coherent state matrix elements⁹. In other words the upper symbol H associated with an operator $\mathcal{H}$ of this type uniquely determines the operator. At the same time the lower symbol h exists for these operators and generally $H(p, q) - h(p, q) = O(\hbar)$, which is unfortunately not easy to see here due to the suppression of $\hbar$ in this work.

The coherent states of the Heisenberg-Weyl group in general, i.e. with an arbitrary fiducial vector, do not (in general) have the property of analyticity¹⁰ but otherwise share all essential properties of the canonical coherent states such as overcompleteness, resolution of unity, universal representation of the operators P and Q by differential operators, etc.

The upper symbol uniquely determines an operator polynomial in P and Q and this is even true for a general operator if the coherent state overlap (reproducing kernel) nowhere vanishes, which happens for uncountably

⁹This is a direct consequence of analyticity most easily exhibited in the complex representation of the canonical coherent states.

¹⁰This is most easily seen in the complex representation of the canonical coherent states.

many fiducial vectors [18]. The existence of the lower symbol is guaranteed for these operators.

An interesting property of the Heisenberg-Weyl group is the relation it establishes between its coherent states and the Wigner/Moyal phase space formulation of quantum mechanics.

For any states $|\psi_1\rangle$ and $|\phi_1\rangle$ the x -representation of $\langle\psi_1|U(p, q)|\phi_1\rangle = \int \psi_1^*(x + q/2)e^{ipx}\phi_1(x - q/2)dx$ is the Fourier transform of a general Wigner function [5][10]. Also, for arbitrary $|\psi_2\rangle$ and $|\phi_2\rangle$,

$$(2\pi)^{-1} \int \langle\psi_1|U(p, q)|\phi_1\rangle \langle\psi_2|U(p, q)|\phi_2\rangle^* dp dq = \langle\psi_1|\phi_1\rangle \langle\psi_2|\phi_2\rangle \quad (16)$$

holds, which essentially is a group orthonormality relation proved by Moyal. But, dropping $\langle\psi_1|$ and $|\psi_2\rangle$, this expression exhibits the resolution of unity in the form $(2\pi)^{-1} \int U(p, q)|\phi_1\rangle \langle\phi_2|U^\dagger(p, q)dp dq = \langle\phi_2|\phi_1\rangle \mathbb{I}$.

Spin coherent states:

The spin coherent states are the group-defined coherent states of the SU(2) group the generators of which satisfy the spin algebra $[S_l, S_m] = i\epsilon_{lmn}S_n$, $l, m, n \in \{1, 2, 3\}$, and ϵ is the totally antisymmetric tensor (Levi-Civita tensor) and summation over repeated indices is understood. One strongly continuous unitary irreducible representation is given by $U(\theta, \phi, \psi) := e^{-i\phi S_3}e^{-i\theta S_2}e^{-i\psi S_3}$, $0 \leq \theta \leq \pi$, $0 \leq \phi < 2\pi$, $0 \leq \psi < 2\pi$, which is called the Euler angle characterization.

The operator U acts on a $(2s + 1)$ -dimensional Hilbert space, where s is the spin. If the fiducial vector is chosen as an eigenstate $|m\rangle$ of S_3 then the operator $e^{-i\psi S_3}$ becomes a simple phase and can be omitted leading to

$$U(\theta, \phi)|m\rangle = e^{-i\phi S_3}e^{-i\theta S_2}|m\rangle = |\theta\phi\rangle \quad (17)$$

closely resembling the case of the canonical coherent states since it involves only two variables.

The expression $\sin\theta d\theta d\phi$ is proportional to the invariant group measure and the group volume is 4π which is finite since the group is compact. Therefore the resolution of unity becomes

$$\mathbb{I} = \frac{2s+1}{4\pi} \int |\theta\phi\rangle \langle\theta\phi| \sin\theta d\theta d\phi \quad (18)$$

and the coherent state overlap reads

$$\begin{aligned}
\langle \theta \phi | \theta' \phi' \rangle &= \langle m | e^{i\theta S_2} e^{i(\phi - \phi') S_3} e^{-i\theta' S_2} | m \rangle \\
&= \sum_{n=-s}^s \langle m | e^{i\theta S_2} | n \rangle \langle n | e^{-i\theta' S_2} | m \rangle e^{i(\phi - \phi') n}
\end{aligned} \tag{19}$$

which is recognized to be expressed in terms of reduced Wigner coefficients of the spin- s representation. The overlap reduces to a simple expression for extremal weight vectors ($m = \pm s$). For $m = s$ it is

$$\begin{aligned}
\langle \theta \phi | \theta' \phi' \rangle &= [\cos(\theta/2) \cos(\theta'/2) e^{i/2(\phi - \phi')}]^{2s} \\
&\quad \times [1 + \tan(\theta/2) \tan(\theta'/2) e^{-i(\phi - \phi')}]^{2s}
\end{aligned} \tag{20}$$

If $m = s$ then every operator $\mathcal{H}$ is uniquely determined by its upper symbol $H(\theta, \phi)$ and thus every operator admits a diagonal representation, i.e. the lower symbol exists [14][24]. This is not necessarily true for a choice of fiducial vector other than the extremal weight ones¹¹.

Observe that the Euler angle characterization is given in non-canonical coordinates. For canonical coordinates, $p = s^{1/2} \sin \theta$ and $q = s^{1/2} \phi$, the overlap becomes

$$\begin{aligned}
\langle pq | p' q' \rangle &= \frac{2s+1}{4\pi s} \left[\frac{1}{2} (1 + s^{-1/2} p)^{1/2} (1 + s^{-1/2} p')^{1/2} \exp\{i s^{-1/2} (q - q')/2\} \right. \\
&\quad \left. + \frac{1}{2} (1 - s^{-1/2} p)^{1/2} (1 - s^{-1/2} p')^{1/2} \exp\{-i s^{-1/2} (q - q')/2\} \right]^{2s}
\end{aligned}$$

and in this notation it can be seen that in the limit $s \rightarrow \infty$ the overlap reduces to the overlap for canonical coherent states, equation (13), for oscillator frequency $\omega = 1$.

Affine coherent states:

The “ax+b”-group, or affine group, is the set $M_+ := \mathbb{R}^+ \times \mathbb{R}$ with the group law $(a', b')(a, b) = (a'a, b' + a'b)$. The group has two nontrivial inequivalent irreducible unitary representations $U_\pm$ with realizations on $L^2(\mathbb{R}^+)$: $[U_\pm(a, b)\psi](x) = a^{1/2} e^{\pm i b x} \psi(ax)$ [26].

Although still referred to as the “ax+b”-group, new coordinates (labels) $a \rightarrow q^{-1}$ and $b \rightarrow p$ will be used throughout this work.

¹¹See [14], page 34, for a counterexample.

The generators of the representations $U_{\pm}$ are the affine kinematical variables Q with $Q > 0$ (position operator) and D (dilation operator) satisfying the affine commutation relation

$$[Q, D] = iQ \quad (21)$$

Then the representations in group coordinates of the second kind are

$$U_{\pm}(p, q) = e^{\pm ipQ} e^{-i \ln q D} \quad (22)$$

Only U_+ will later be important, so $U := U_+$.

As in the canonical and spin case, the main interest will concentrate on the group action on extremal weight vectors. The minimum uncertainty states, satisfying the Heisenberg uncertainty equation $\Delta Q \Delta P = \langle Q \rangle / 2$, are a 2-parameter family given in x -representation as [27] $\eta_{\alpha, \beta}(x) = N_{\alpha, \beta} x^{\alpha} e^{-\beta x}$.

Setting the mean of Q to 1 will lead to a 1-parameter family of minimum uncertainty states

$$\eta_{\beta}(x) = N_{\beta} x^{\beta-1/2} e^{-\beta x} \quad (23)$$

with the normalization $N_{\beta} = (2\beta)^{\beta} \Gamma^{-1/2}(2\beta)$.

The vectors $|\eta_{\beta}\rangle$ satisfy $(Q - 1 + i\beta^{-1}D)|\eta_{\beta}\rangle = 0$, which is the analog of $(\omega^{1/2}Q + i\omega^{-1/2}P)|0\rangle = 0$ in the canonical case.

This again leads to a complex polarization condition (as in the canonical case) and to analytic properties of the affine coherent states (most easily recognized in the overlap).

The affine coherent states are defined as

$$|pq\rangle = U(p, q)|\eta_{\beta}\rangle \quad (24)$$

and the β -dependence will often be left implicit, i.e. $|\eta_{\beta}\rangle =: |\eta\rangle$.

For the whole parameter range $0 < \beta$ the overlap is

$$\langle pq|rs\rangle = (qs)^{-\beta} 2^{-2\beta} [(q^{-1} + s^{-1}) + i\beta^{-1}(p - r)]^{-2\beta} \quad (25)$$

The left-invariant group measure is proportional to $dp dq$ whereas the right-invariant group measure is proportional to $dp dq/p$. This is the first example where the two measures differ and it follows from the general discussion that square integrability of the representation, necessary for a resolution of unity, can impose restrictions on the choice of fiducial vectors.

This indeed happens, and there is a fiducial vector admissibility condition [25][26][27]

$$\langle Q^{-1} \rangle = \int_0^\infty x^{-1} |\eta_\beta(x)|^2 dx < \infty \quad (26)$$

This condition is violated if $0 < \beta \leq 1/2$ and fulfilled for $\beta > 1/2$ in which case the resolution of unity reads

$$\mathbb{1} = (1 - \frac{1}{2\beta})(2\pi)^{-1} \int |pq\rangle \langle pq| dp dq \quad (27)$$

Observe that adding the phase factor $\exp\{-i\beta^{1/2}(p-r)\}$ to the overlap and translating q to $q + \beta^{1/2}$ one obtains in the limit $\beta \rightarrow \infty$ the overlap of the canonical coherent states (again for unit angular frequency) [28].

2.4 The standard coherent state path integral

Let $\mathcal{H}$ be the Hamiltonian operator for a quantum mechanical system which is supposed not to depend on time explicitly¹². Then the time evolution operator is $\hat{U}(t'' - t') = \exp\{-i\mathcal{H}(t'' - t')\}$ and w.l.o.g. $t' = 0$, $t'' = T$. Then $|\psi(t)\rangle = \hat{U}(t)|\psi(0)\rangle$ solves the Schrödinger equation.

The standard approach to a path integral is the factorization of the evolution operator $\exp\{-i\mathcal{H}T\}$ into N equal terms $\exp\{-i\mathcal{H}\epsilon\} \dots \exp\{-i\mathcal{H}\epsilon\}$ with $N\epsilon = T$. Then resolutions of unity are inserted between these factors, in this case given by the expression $\mathbb{1} = \int |l\rangle \langle l| \delta l$ belonging the coherent states labeled by l . This leads to

$$\langle l''; T | l'; 0 \rangle \equiv \int \prod_{k=0}^N \langle l_{k+1} | e^{-i\epsilon\mathcal{H}} | l_k \rangle \prod_{k=1}^N \delta l$$

and since this is an identity it holds in the limit $\epsilon \rightarrow 0$ taken as a final step. As usual $l_{N+1} := l''$ and $l_0 := l'$. For small ϵ the approximation for each factor

$$\begin{aligned} \langle l_{k+1} | e^{-i\epsilon\mathcal{H}} | l_k \rangle &\approx \langle l_{k+1} | (1 - i\epsilon\mathcal{H}) | l_k \rangle \\ &= \langle l_{k+1} | l_k \rangle [1 - i\epsilon \langle l_{k+1} | \mathcal{H} | l_k \rangle \langle l_{k+1} | l_k \rangle^{-1}] \\ &\approx \langle l_{k+1} | l_k \rangle \exp\{-i\epsilon \langle l_{k+1} | \mathcal{H} | l_k \rangle \langle l_{k+1} | l_k \rangle^{-1}\} \end{aligned} \quad (28)$$

¹²Generalization to time-dependent Hamiltonians is straightforward and will involve the time-ordering operator $\mathbf{T}$.

Canonical coherent states	
generators	(P, Q)
phase space variables	(p, q)
coherent states	$ pq\rangle = \exp\{-iqP\} \exp\{ipQ\} 0\rangle$
overlap	$\langle pq p'q'\rangle = e^{i/2(p'q - q'p)} e^{-1/4[(p-p')^2 + (q-q')^2]}$
canonical 1-form	$i\langle pq d pq\rangle = p dq$
induced metric	$d\sigma^2 = 2[\ d pq\rangle\ ^2 - \langle pq d pq\rangle ^2] = dp^2 + dq^2$
Spin coherent states	
generators	(S_2, S_3)
phase space variables	(θ, ϕ) or (p, q)
coherent states	$ \theta\phi\rangle = e^{-i\phi S_3} e^{-i\theta S_2} s\rangle$
overlap (non-can. coord.)	$\langle \theta\phi \theta'\phi'\rangle = \sum_{n=-s}^s \langle m e^{i\theta S_2} n\rangle \langle n e^{-i\theta' S_2} m\rangle e^{i(\phi-\phi')n}$
overlap (can. coord.)	$\langle pq p'q'\rangle = \frac{2s+1}{4\pi s} [\frac{1}{2}(1+s^{-1/2}p)^{1/2}(1+s^{-1/2}p')^{1/2} \times \exp\{\frac{i}{2}s^{-1/2}(q-q')\} + \frac{1}{2}(1-s^{-1/2}p)^{1/2}(1-s^{-1/2}p')^{1/2} \times \exp\{-\frac{i}{2}s^{-1/2}(q-q')\}]^{2s}$
non-canonical 1-form	$i\langle \theta\phi d \theta'\phi'\rangle = s \sin \theta d\theta d\phi$
canonical 1-form	$i\langle pq d pq\rangle = p dq$
metric (non-canonical)	$d\sigma^2 = s(d\theta^2 + \sin^2 \theta d\phi^2)$
metric (canonical)	$d\sigma^2 = (1-p^2/s)^{-1} dp^2 + (1-p^2/s) dq^2$
Affine coherent states	
generators	(Q, D)
phase space variables	(p, q)
coherent states	$ pq\rangle = \exp\{ipQ\} \exp\{-i \ln q D\} \eta\rangle$
overlap	$\langle pq rs\rangle = (qs)^{-\beta} 2^{-2\beta} [(q^{-1} + s^{-1}) + i\beta^{-1}(p-r)]^{-2\beta}$
canonical 1-form	$i\langle pq d pq\rangle = p dq$
induced metric	$d\sigma^2 = 2[\ d pq\rangle\ ^2 - \langle pq d pq\rangle ^2] = \beta^{-1} q^2 dp^2 + \beta q^{-2} dq^2$

Table 1: Coherent states

is correct to lowest order in ϵ [22]. With $H(l_{k+1}; l_k) := \langle l_{k+1}|l_k\rangle^{-1} \langle l_{k+1}|\mathcal{H}|l_k\rangle$ it follows that

$$\langle l''; T|l'; 0\rangle = \lim_{\epsilon \rightarrow 0} \int \prod_{k=0}^N [\langle l_{k+1}|l_k\rangle \exp\{-i\epsilon H(l_{k+1}; l_k)\}] \prod_{k=1}^N \delta l \quad (29)$$

This is a well-defined expression if the integrals converge¹³. It is customary at this point to interchange the integrations and the limit although this is

¹³If the integrals do not converge, the last approximation in Eq. (28) - although correct in the integrand - simply did not result in an integrable function and hence can not be made.

not justified and replace the variables l_k by real functions $l(t)$. With

$$\prod_{k=0}^N \langle l_{k+1} | l_k \rangle = \prod_{k=0}^N [1 - \langle l_{k+1} | (|l_{k+1}\rangle - |l_k\rangle)] \approx \exp\left\{-\sum_{k=0}^N \langle l_{k+1} | (|l_{k+1}\rangle - |l_k\rangle)\right\}$$

valid to first order in the difference vectors the (formal) path integral is given by

$$\langle l''; T | l'; 0 \rangle = \mathcal{N} \int \exp\{i \int [i \langle l(t) | \dot{l}(t) \rangle - H(l)] dt\} \mathcal{D}l \quad (30)$$

where $H(l) = \langle l(t) | \mathcal{H} | l(t) \rangle$ is the upper symbol, $\mathcal{D}l = \lim_{\epsilon \rightarrow 0} \prod_{k=1}^N \delta l_k$ and $\mathcal{N}$ is a (formal, possibly infinite, possibly zero) normalization constant.

There is a second derivation of a coherent state path integral [14], this time involving the lower symbol $h(l)$ which is assumed to exist for the Hamiltonian $\mathcal{H}$ under consideration. Given some restrictions on $h(l)$ ¹⁴ the operator $F(t) := \int e^{-ith(l)} |l\rangle \langle l| \delta l$ is bounded by unity and the Cauchy-Schwarz inequality gives $|\langle \phi | F(t) | \psi \rangle| \leq \| |\psi\rangle \| \cdot \| |\phi\rangle \|$ for all $|\phi\rangle$ and $|\psi\rangle$. The strong limit $s - \lim_{t \rightarrow 0} [\mathbb{1} - F(t)]/(it)$ exists and is $\mathcal{H}$ on the coherent states. This is enough [21] to ensure $s - \lim_{N \rightarrow \infty} [F(T/N)]^N = \exp\{-iT\mathcal{H}\}$ on the whole Hilbert space $\mathfrak{H}$. Application of this, and setting $\epsilon := T/N$, $l_{N+1} := l''$, $l_0 := l'$, leads to

$$\langle l'' | \exp\{-iT\mathcal{H}\} | l' \rangle = \lim_{N \rightarrow \infty} \int \prod_{n=0}^N \langle l_{n+1} | l_n \rangle \prod_{n=1}^N \exp\{-i\epsilon h(l_n)\} \delta l_n \quad (31)$$

which is a valid (discrete) “path integral”.

Interchange of the integrations and the limit results in an expression

$$\langle l'' | \exp\{-iT\mathcal{H}\} | l' \rangle = \int \exp\{i \int [i \langle l(t) | \dot{l}(t) \rangle - h(l)] dt\} \mathcal{D}l \quad (32)$$

for the coherent state propagator which is different from equation (30) since in general $H(l) \neq h(l)$ [compare the definitions of the upper and lower symbol, equations (9), (10)].

This difference clearly shows the formal character of the standard path integral since it suggests the equality of two expressions which obviously are not equal. The discrete lattice formulations, on the other hand, are rigorously derived and valid.

¹⁴namely $\int |h(l_1)h(l_2)\langle l|l_1\rangle\langle l_1|l_2\rangle\langle l_2|l|\delta l_1\delta l_2| < \infty$

One remark about the connection of classical and quantum mechanics is in place here. Consider the quantum action functional $I = \int \langle \psi | i\partial_t - \mathcal{H} | \psi \rangle dt$ from which the Schrödinger equation $i\dot{|\psi\rangle} = \mathcal{H}|\psi\rangle$ can be derived by requesting stationarity under unrestricted variations of the vector $|\psi(t)\rangle$. W.l.o.g the variation can be carried out over unit vectors only, also excluding mere phase variations $|\psi_t\rangle = \exp\{i\alpha(t)\}|\psi\rangle$ [22].

The action in the exponent of the path integral, $I = \int [i\langle l(t) | \dot{l}(t) \rangle - H(l)] dt$, is the classical action if the upper symbol is regarded as the classical Hamiltonian according to the weak correspondence principle. Variation will lead to the classical equations of motion. This classical action functional is identical in form with the quantum action functional, the only difference being that the variations range only over the special set of coherent state unit vectors instead of over all unit vectors. Regarded in this light, and referring to an analog with thermodynamic equilibrium, “quantum mechanics is just classical mechanics with all stops removed” [22]

2.5 Mathematical aspects of path integration

In the last section it was already stressed that the path integral, as it stands, is only a formal expression. The same (coherent state) propagator was expressed in two different path integrals. And it gets worse. The same path integral expression can be shown to stem from completely different propagators.

To realize this, consider Eq. (30) and let the arbitrary coherent states $|l\rangle$ now be the coherent states of the Heisenberg-Weyl group $|pq\rangle$ which are supposed to be centered, i.e. $\langle pq | Q | pq \rangle = 0$ and $\langle pq | P | pq \rangle = 0$. These could be the canonical coherent states. Because of the translation property Eq. (14), one has $id|pq\rangle = id(e^{-iqP} e^{ipQ} |\psi_0\rangle) = [-Qdp + (P + p)dq]|pq\rangle$, and hence $\langle pq | i\partial_t | pq \rangle = p\dot{q}$ holds.

The action functional reads $I = \int [p\dot{q} - H(p, q)] dt$ and the path integral is now

$$\mathcal{N} \int \exp\{i \int [p\dot{q} - H(p, q)] dt\} \mathcal{D}p \mathcal{D}q \quad (33)$$

However, this is the very same expression one obtains for the conventional phase space path integral even though the propagators are not the same. The coherent state propagator $\langle p''q''; T | p'q'; 0 \rangle$ is certainly not equal to $\langle q''; T | q'; 0 \rangle$. Even the interpretation of “ $\mathcal{D}p$ ” and “ $\mathcal{D}q$ ” is different. In the conventional path integral the “ p ’s” are integrated out (“there is always one more dp than dq ”) whereas in the coherent state interpretation there are

the same number of p - and q -integrations¹⁵. This does not mean that the expressions are simply wrong since it was said before that the path integral at this level is merely formal. The last rigorous expressions, at the “lattice level”, are quite different. For the coherent state propagator this is Eq. (29), now with p and q as labels:

$$\begin{aligned} \langle p'' q''; T | p' q'; 0 \rangle &= \lim_{\epsilon \rightarrow 0} \int \prod_{k=0}^N \left[\langle p_{k+1} q_{k+1} | p_k q_k \rangle \right. \\ &\quad \left. \times \exp \{ -i\epsilon H(p_{k+1}, q_{k+1}; p_k, q_k) \} \right] \prod_{k=1}^N \frac{dp_k dq_k}{2\pi} \quad (34) \end{aligned}$$

The conventional propagator, on the other hand, is expressed as

$$\begin{aligned} \langle q''; T | q'; 0 \rangle &= \lim_{N \rightarrow \infty} (2\pi)^{-N+1} \int \exp \left\{ i \sum_{l=0}^N \left[p_{l+1/2} (q_{l+1} - q_l) \right. \right. \\ &\quad \left. \left. - \epsilon H(p_{l+1/2}, \tfrac{1}{2}(q_{l+1} + q_l)) \right] \right\} \prod_{l=0}^N dp_{l+1/2} \prod_{l=1}^N dq_l \end{aligned}$$

where $q_{N+1} = q''$ and $q_0 = q'$ and the indices $l + 1/2$ indicate that the resolutions of unity for the p -representation have been inserted at different (intermediate) times. Otherwise the Heisenberg uncertainty principle would be violated. This is because p and q are “sharp” eigenstates in their own representations as opposed to the coherent state representation, where they are both mean values which can be specified at the same time.

Consider now the path integral $\mathcal{N} \int \exp \{ i \int [p \dot{q} - H(p, q)] dt \} \mathcal{D}p \mathcal{D}q$ in its own right, i.e. no matter what propagator it was derived from. What can be said about this expression?

First, note that the functional “measures” $\mathcal{D}p$ and $\mathcal{D}q$ are modeled in a way to be an analog of the homogeneous, translation invariant Lebesgue measure. But it is known [19][20] that such a measure does not exist. It lacks the property of countable additivity (it is only finitely additive). Hence, the so-called path integral is not an integral but a linear functional.

Second, even if the measure existed, the integral would be improper since it is of the form $\mathcal{N} \int \exp \{ iI \} \mathcal{D}p \mathcal{D}q$ which is not absolutely integrable. So some sort of regularization is necessary to remove ambiguities that go along with this fact.

The Feynman path integral (the conventional real time path integral) has been constructed to be a formulation of quantum mechanics equivalent

¹⁵This has deep consequences concerning the geometric nature of the formulation. See section (2.6).

to the Schrödinger formulation [36]. It was pointed out in the very beginning of the present work that the Schrödinger formulation resides in Cartesian coordinates only. Not surprisingly, the Feynman path integral has the same limitations. Superficially, the path integral $\mathcal{N} \int \exp\{i \int [p\dot{q} - H(p, q)] dt\} \mathcal{D}p \mathcal{D}q$ looks covariant under canonical transformations, but it is not. On the “lattice level”, the last rigorous formulation defining the formal path integral, there is no such covariance. Naturally, canonical transformations can be performed, but the path integral has to be rederived in every case and will generally look quite different then¹⁶.

The coherent states on the other hand merely change labels under canonical transformations (apart from phase factors). Does that mean that the latter can be performed in the path integrand? Certainly not. The same things said above about the Feynman path integral hold true here as well. The standard coherent state path integral is defined by the lattice approach and there is no covariance under canonical transformations.

A last remark about the formal path integral: It is well-known that operator ordering ambiguities arise when, according to the “royal route”, i.e. Schrödinger quantization, classical variables are promoted to operators. Where is this ordering ambiguity in the path integral? It is concealed in the formal expression but clearly present on the lattice level. There, an expression like $q_k^2 p_{k-1/2} q_{k-1}$ would correspond to an operator $Q^2 P Q$ whereas $q_k^3 p_{k-1/2}$ would correspond to $Q^3 P$ etc.

It might be unjust to say the path integral hides more truth than it exhibits since it gives a certain amount of intuitive insight and raises many interesting questions. However, it would be foolish to try to rigorously derive anything from an expression that is so ill-defined. Questions like “what is the nature of the paths” can be answered heuristically but are, strictly speaking, meaningless ([14], pages 67-68).

2.6 Mathematical tools

How to overcome all the aforementioned ambiguities will be the content of section (2.7). The formulation of a well-defined coherent state path integral will require some mathematical background not previously encountered in standard path integration. This section introduces these mathematical tools. For simplicity everything will be formulated for a single degree of freedom.

¹⁶See e.g. [30], page 63-66 for the path integral in spherical coordinates, or page 67-78 for the path integral in general coordinates.

The “shadow metric”

Classical mechanics is described by Hamilton’s equations of motion

$$\dot{q} = \partial_p h(p, q) \quad \dot{p} = -\partial_q h(p, q) \quad (35)$$

where $h(p, q)$ is the classical Hamiltonian. These equations are derived from an equivalence class of action functionals, $I = \int pdq + dF_0 - hdt$, by requirement of stationarity under variations which hold both endpoints of phase-space paths fixed. Thus the C^1 function F makes no contribution to the equations of motion.

Note that Hamilton’s equations are often derived from an action $S = \int [p\dot{q} - H(p, q)]dt$, holding only the position q fixed at the end-points. The reason for this is that in the standard path integral holding both endpoints fixed would lead to an overspecification since the p ’s are completely integrated out (remember: “there is always one more dp than dq ”). Certainly, this can not be a geometric formulation when the phase space variables are treated on an unequal footing. In this approach, Hamilton’s equations can not be derived from an action $S = \int [-q\dot{p} - H(p, q)]dt$ without first changing the problem by partial integration to the standard action given above. The problem can not be overcome with the traditional coherent state path integral since the path “integral” is ill-defined as it stands. But, if there is a sort of well-defined coherent state path integral, then the fact that p and q are treated in the same way promises the reward of a geometric nature of quantization.

The equations of motion are covariant under canonical (C^1 coordinate) transformations [31], which imply the existence of an underlying geometrical structure [32]: Phase space is a differentiable manifold M endowed with a nondegenerate, closed, symplectic two-form ω . So locally $\omega = d(\theta + dF_0)$, and the action functional is $I = \int (\theta + dF_0 - hdt)$. By Darboux’s theorem local coordinates can be introduced such that $\theta = pdq$, $F_0 = F_0(p, q)$ and $\omega = dp \wedge dq$ leading to the equations of motion (35).

Covariance under canonical (coordinate) transformations implies on the other hand that coordinatized expressions, e.g. for the Hamiltonian, have no definite physical meaning by themselves. Consider a Hamiltonian of the form $h(p, q) = p^2 + q^2$ which is supposed to correspond physically to the harmonic oscillator. In a different coordinate system (here: polar coordinates in phase space) the same Hamiltonian becomes $\bar{h}(\bar{p}, \bar{q}) = \bar{p}$. Without further specifications each of these expressions would not be connected to any particular physical system.

On the other hand, if the phase space (M, ω) is augmented by a Riemannian metric $d\sigma^2$ then mathematical expressions like $h(p, q)$ gain a unique physical meaning. Consider the example above: $(M, \omega, d\sigma)$ with metric $d\sigma^2 = dp^2 + dq^2$ (Cartesian metric) would show that $h(p, q) = p^2 + q^2$ is the harmonic oscillator Hamiltonian whereas $h(p, q) = p$ clearly is not. On the other hand choosing the polar metric $d\sigma^2 = (2p)^{-1}dp^2 + (2p)dq^2$ would imply that $h(p, q) = p$ is the harmonic oscillator Hamiltonian and $h(p, q) = p^2 + q^2$ would correspond to something more exotic.

The metric $d\sigma^2$ is often called a “shadow” or “secret” metric [33].

Brownian motion

A shadow metric appended to phase space has another effect, namely it supports Brownian motion (sometimes called the Wiener process) on the Riemannian phase space manifold. The symplectic structure alone was not enough since only with a metric can one define the Laplace-Beltrami operator

$$\Delta_{LB} := \sqrt{g}^{-1} \partial_a \sqrt{g} g^{ab} \partial_b \quad (36)$$

where g^{ab} , $a, b \in \{p, q\}$, is the metric tensor ($d\sigma^2 = g^{ab} dx_a dx_b$) and g is its determinant. Let $K(p'', q'', p', q', t) := [\exp\{t\Delta_{LB}\}](p, q, p', q')$ be the heat kernel, i.e. the solution to the diffusion equation

$$\partial_t K = \frac{1}{2} \nu \Delta_{LB} K \quad (37)$$

with initial condition $K(p, q, p', q'; 0) = \delta(p - p')\delta(q - q')$. The parameter ν is called the diffusion constant. Then

$$\int d\mu_{W;p'',q'',p',q'}^{\nu;T} := K(p'', q'', p', q'; \nu T) \quad (38)$$

unambiguously defines a measure pinned at (p', q') for $t = 0$ and at (p'', q'') for $t = T$ [26]. It is called the pinned or conditional[29] Wiener measure with diffusion constant ν and is a true measure on path space concentrated on continuous but nowhere differentiable paths. Often the parameters p'', q'', p' and q' will be suppressed or - in a slight abuse of notation - be written on the integral sign as upper and lower limits of the integration.

So the paths are continuum random walks or, better, stochastic processes (See [29] for an elementary introduction or [34] for a more mathematical approach). To be more precise they are so-called Brownian bridges and are




Figure 1: Brownian Paths

intimately related to Brownian motion which is the key to understand the nature of the paths¹⁷.

Standard Brownian motion (or the Wiener process) is the family $(X_t)_{0 \leq t}$ of centered, i.e. mean zero, Gaussian random variables with covariance $\langle X_t X_s \rangle = \min(t, s)$. Here $\langle \cdot \rangle$ denotes the expectation value. Since the process is Gaussian, the first two moments specify all higher moments. This fact can be expressed in the form of the moment-generating functional $\langle e^{ibX} \rangle = e^{-b^2 \langle X^2 \rangle / 2}$.

It was already stated that the Brownian paths are continuous but nowhere differentiable. This can be seen as follows:

From the generating functional the second and fourth moment are readily computed as $\langle (X_t - X_s)^2 \rangle = |t - s|$ and $\langle (X_t - X_s)^4 \rangle = 3|t - s|^2$. Consider now $\langle (X_{t+\Delta} - X_t)^2 \rangle = \Delta$ and write it as $\Delta = A\Delta + B\Delta$. The first contribution comes from continuous paths and here Δ can be interpreted as arising from the slope of the path on the interval $[t, t + \Delta]$. The second contribution comes from discontinuous paths and in this case Δ can be seen as being proportional to the capturing probability for such a path (see figure 1). The aim is to show that there are no contributions from discontinuous paths, i.e. $B = 0$. Continuing the argumentation with the Δ -parts coming from continuous/discontinuous paths one would expect an expression $\langle (X_{t+\Delta} - X_t)^4 \rangle = A'\Delta^2 + B'\Delta$ since in the continuous part the slope would be squared now whereas the capturing probability stays the same, i.e. linear in Δ . But

¹⁷Brownian processes enter the picture naturally since they are connected to random walks with diffusion, which in turn are connected to the Laplace-Beltrami operator by the diffusion equation.

the fourth moment really is $\langle (X_{t+\Delta} - X_t)^4 \rangle = 3\Delta^2$ showing that there is no contribution from discontinuous paths (and hence the same is true for the second moment as well, i.e. $B = 0$). The conclusion is that Brownian paths are continuous with probability one.

The statement about continuity can be proved in a less pictorial way by recognizing that the fourth moment fulfills the properties required for a theorem of Kolmogorov [35] which says: If there are $\alpha > 0$, $\beta > 0$, $C > 0$ such that $\langle |Z_{t_2} - Z_{t_1}|^\alpha \rangle \leq C|t_2 - t_1|^{1+\beta}$ for all t_1, t_2 then the process Z is continuous with probability one.

To show that Brownian paths are almost nowhere differentiable, the moment generating functional $\langle e^{itX} \rangle$ is integrated over t in the following way: $\int e^{-t^2/2} \langle e^{itX} \rangle dt = \int e^{-t^2/2} e^{-t^2 \langle X^2 \rangle / 2} dt \propto (1 + \langle X^2 \rangle)^{-1/2} = \langle e^{-X^2/2} \rangle$. These relations hold since X is Gaussian. Consider the stochastic process $Y := \frac{X(t) - X(s)}{|t-s|^\beta}$ which becomes the derivative of X for $\beta = 1$ and $t \rightarrow s$. Then $\langle \exp\{-\frac{1}{2} \frac{[X(t) - X(s)]^2}{|t-s|^{2\beta}}\} \rangle = (1 + |t-s|^{1-2\beta})^{-1/2}$. For $\beta = 1$ and $t \rightarrow s$ this assumes the form $\langle e^{-[\dot{X}^2(t)]/2} \rangle = 0$ which says that $\dot{X}$ is infinity almost everywhere. Thus Brownian paths are almost nowhere differentiable.

General Brownian motion is connected to standard Brownian motion by $B_t = \sqrt{\nu}X_t + c$ where the offset c is a constant and so is ν (often called the diffusion constant).

If X_t is standard Brownian motion with $X_0 = 0$, then scaling shows that $X_t^* := l^{-1}(X_{s+tl^2} - X_s)$ is Brownian motion again with $X_0^* = 0$. The standard Brownian bridge is defined as the stochastic process

$$\bar{X}_t := X_t^* - tX_1^* \quad (39)$$

and has mean zero and covariance¹⁸

$$\langle \bar{X}_t \bar{X}_s \rangle = \min(t, s) - ts \quad (40)$$

It follows that $\bar{X}_0 = \bar{X}_1 = 0$, and this is why the standard Brownian bridge is sometimes called “circular” Brownian motion.

The general Brownian bridge is connected to the standard one first by rescaling of the time (to get an arbitrary time-interval instead of $[0, 1]$): $\tilde{X}_t := X_t^* - \frac{t}{T}X_T^*$ and hence $\langle \tilde{X}_t \tilde{X}_s \rangle = \min(t, s) - ts/T$. Then the deterministic classical path with initial point x' at time 0 and final point x'' at time T is added leading to the expression $\bar{B}_t = x'(T-t)/T + x''t/T + \tilde{X}_t$.

¹⁸The Brownian bridge can also be defined as the centered Gaussian process with this covariance. Then the other properties follow, see e.g. [34].

In the path integral the paths will be restricted by initial and final conditions. This is exactly why the Brownian bridge rather than Brownian motion will be relevant.

Stochastic integrals

Let X be Brownian motion. It was stated above that Brownian paths are nowhere differentiable. But since $\partial_t \partial_s \min(s, t) = \delta(t - s)$ one can talk about a formal derivative $W_t = \dot{X}_t$ with covariance $\langle W_s W_t \rangle = \delta(t - s)$ termed “white noise”. This distribution, called a generalized stochastic process, can produce bona-fide stochastic variables when integrated against suitable test functions. Let f be such a function then

$$W(f) := \int_0^\infty f(t) W_t dt =: \int_0^\infty f(t) dX_t \quad (41)$$

is well-defined. The last expression can also be understood in another way. The problem is that dX_t is of unbounded variation on any time interval and does not represent a true measure. But by partial integration one obtains $\int \dot{f} X_t dt$ which can be interpreted as a Lebesgue-Stieltjes integral (provided integrability conditions on $\dot{f}$ hold).

Let Y be another Brownian motion, possibly independent of X . The integral

$$\int X dY \quad (42)$$

can not be given meaning by partial integration (dX would be just as unsuitable a measure as dY). One has to adopt a rule on how to define this expression. The most common rules are the Itô and Stratonovich interpretations. Both rules define the integral as the continuum limit of discrete random walks on a lattice. The Itô rule says $\int X dY := \lim \sum X_l (Y_{l+1} - Y_l)$ whereas Stratonovich uses the “midpoint rule”

$$\int X dY := \lim \sum \frac{1}{2} (X_{l+1} + X_l) (Y_{l+1} - Y_l) \quad (43)$$

Because of the unbounded variations of the Brownian paths these two expressions are generally different (as opposed to a “normal” deterministic integral, where they would coincide).

The Itô rule is favored by most mathematicians since it has theoretical advantages which are based on the independent increments of a Brownian process [29]. This leads directly to the famous Itô calculus (see e.g. [34]),

a stochastic differential calculus. One of its rules says that the square of the stochastic differential is deterministic, namely $dX_t^2 = dt$. This is more than saying that the mean of this squared differential is dt , which was already recognized by Feynman in his original work on path integration [36]: “Although the average value of the displacement of a particle in the time dt is vdt , where v is the mean velocity, the mean of the square of this displacement is not of order dt^2 , but only of order dt ”. An application of this rule on the metric $d\sigma^2 = dx_a g^{ab} dx_b$ shows that, as a stochastic variable, $d\sigma^2 = 2\nu dt$, where ν is the diffusion constant and the factor 2 indicates the dimensionality of the phase space manifold [37].

On the other hand the Stratonovich rule has the advantage that common rules of calculus still apply (which is generally not true for the Itô-integral, see e.g. [34]). Canonical transformations can be performed on stochastic processes in formally the same way as for deterministic functions when the Stratonovich rule is applied.

For more information on Brownian motion, other stochastic processes and stochastic calculus and integration see e.g. [29][34][35].

Functional integration with Wiener measure

A standard way to get a well-defined path integral is to analytically continue the time-evolution operator to imaginary time (Wick rotation), thus obtaining the operator $\exp\{-TH\}$ characteristic for the diffusion equation. Here H is the Hamilton operator and if it is of the form $H = H_0 + V$, where $H_0 = -\frac{1}{2}\Delta$ is the Hamiltonian for a free particle with unit mass, one is led to the Euclidean configuration space path integral [29][30]

$$\begin{aligned} K_E(x'', x'; T) &= \langle x'' | \exp\{-T[H_0 + V(x)]\} | x' \rangle \\ &= \lim_{N \rightarrow \infty} (2\pi\epsilon)^{-N/2} \prod_{k=1}^{N-1} \int dx_k \exp\left\{-\sum_{j=0}^{N-1} \left[\frac{1}{2\epsilon}(x_{j+1} - x_j)^2 + \epsilon V(x_j)\right]\right\} \\ &=: \int \exp\left\{-\int \left[\frac{1}{2}\dot{x}^2 + V(x)\right]dt\right\} \mathcal{D}x =: \int \exp\left\{-\int V dt\right\} d\mu_W \end{aligned}$$

where $d\mu_W = \exp\{-\int \frac{1}{2}\dot{x}^2\} \mathcal{D}x$ is a Wiener measure pinned at the endpoints.

This result is known as the Feynman-Kac formula and one may speak of functional integration to emphasize its well-definedness. Observe, however, that the Wiener measure arises only because of the special form of the Hamiltonian as a free Hamiltonian plus a potential. This approach has

nothing to do with the coherent state path integral of the next section except that it uses the same mathematical tools.

For further studies, e.g. for a connection between such path integrals and stochastic expectations, see e.g. [29].

Improper integrals

The main problems with the path integral were the ill-definedness of the measure and the ambiguity arising from the fact that, even with a true measure, the path integral in its standard form would be an improper integral of the form $\int \exp\{iI\} \mathcal{D}l$.

To give a well-defined meaning to an improper integral one has to adopt a particular regularization. Consider the toy example

$$\int_{-\infty}^{\infty} e^{iy^2/2} dy := \lim_{\nu \rightarrow \infty} \int_{-\infty}^{\infty} e^{iy^2/2 - y^2/(2\nu)} dy = \sqrt{2\pi i} \quad (44)$$

Here a regularization factor was introduced which rendered the integrand absolutely integrable. The improper integral was defined by this regularization. Although the result $\sqrt{2\pi i}$ seems natural (in the sense that if one simply denied the problem and integrated $e^{-y^2/(2i)}$ according to the usual rule for Gaussian integrals one would be led to the very same answer), it must be stressed that different regularizations could very well lead to different results.

And different results for different regularizations are exactly what will be encountered in the path integral in the next section. The idea of a continuous regularization (or convergence) factor should be kept in mind.

2.7 The coherent state path integral with Wiener measure

Consider again the expression for the coherent state path integral [Eq. (30)]:

$$\langle l''; T | l'; 0 \rangle = \mathcal{N} \int \exp\{i \int [i \langle \dot{l}(t) | l(t) \rangle - H(l)] dt\} \mathcal{D}l$$

where $H(l)$ is again the upper symbol. When trying to apply the stationary phase approximation to the action in the integrand, it turns out that the resulting first-order extremal equations are generally incompatible with the boundary conditions $l'' = l(T)$ and $l' = l(0)$ [38] [22]. The idea is to introduce an extra term $\frac{1}{2}i\epsilon \langle \dot{l} | (1 - |l\rangle\langle l|) | \dot{l} \rangle$ into the action which changes the extremal equations to second order, compatible with the boundary conditions, and taking the limit $\epsilon \rightarrow 0$ as a final step. The fact that this

term is chosen and no other has a reason which will be clarified with the example of the canonical coherent states. The coherent state overlap for a short time interval (and with the notations from the path integral) is $\langle p_{l+1}q_{l+1}|p_lq_l\rangle = \exp\{i(\frac{p_{l+1}+p_l}{2})(q_{l+1}-q_l) - \frac{\epsilon}{4}[\frac{p_{l+1}-p_l}{\epsilon^2} + \frac{q_{l+1}-q_l}{\epsilon^2}]\epsilon\}$. In the limit $N \rightarrow \infty$, i.e. $\epsilon \rightarrow 0$ the second term in the exponent vanishes. But letting one ϵ “by force” not become an infinitesimal dt the term turns out to be the regularization factor $e^{-\frac{\epsilon}{4}\int(\frac{dp^2}{dt^2} + \frac{dq^2}{dt^2})dt}$ introduced above (here for the canonical coherent states). At this level it is “bad mathematics” to “save” one ϵ but it works! Later this procedure will be put on a firm mathematical ground.

But one has to point out that the correctness of this procedure can only be shown a posteriori which means after evaluating the stationary phase method for the well-defined lattice expressions and comparison with the results obtained from this continuous selection method [22].

Assuming correctness (which was shown for the canonical and spin coherent states in [38]) the path integral assumes the form

$$\begin{aligned} & \langle l''; T | l'; 0 \rangle \\ &= \lim_{\epsilon \rightarrow 0} \mathcal{N} \int \exp\{i \int [i \langle l | \dot{l} \rangle + \frac{1}{2} i \epsilon \langle \dot{l} | (1 - |l\rangle \langle l|) | \dot{l} \rangle - H(l)] dt\} \mathcal{D}l \end{aligned} \quad (45)$$

Note that in the exponent two terms of clear geometric character appear: First, $i \langle l | d | l \rangle$ is recognized to be the canonical one-form θ [see the discussion in 2.6], which for the canonical coherent states becomes $\theta = pdq$. Second, $\frac{1}{2}(\langle l | d \rangle (1 - |l\rangle \langle l|) (d | l \rangle))$ is a Riemannian metric $d\sigma^2$, which for the canonical coherent states becomes $d\sigma^2 = dp^2 + dq^2$.

Here appears for the first time a new structure, a metric, which in this case is induced by the coherent states themselves. For the canonical coherent states this turns out to be the Cartesian metric, so there is reason to believe that they are in some way associated with a flat phase-space. This point will be discussed later in a more general context.

The things to be learned from Eq. (45) and to be kept in mind are

- 1) the association of a metric structure with the phase space
- 2) a continuous regularization in the path integral (as opposed to the usual time-slicing and discretization)

Now the continuous regularization of the path integral by a factor including a metric on phase space is taken as a starting point. To begin with,

the metrics under consideration will be the flat, spherical or hyperbolic metric. Later, generalizations to arbitrary metrics on 2-dimensional phase space manifolds without symmetry will be discussed. Instead of the abstract notation $i\langle l|\dot{l}\rangle$ and $\frac{1}{2}i\epsilon\langle\dot{l}|(1-|l\rangle\langle l|)|\dot{l}\rangle$ for the canonical one-form and the metric respectively, canonical coordinates will be used, i.e. those coordinates for which the canonical one-form is pdq .

In the following path integral the total time-derivative of an arbitrary function $G(p, q)$ is included which, for a simply connected phase space manifold, will simply be an unimportant overall phase factor¹⁹. The classical Hamiltonian is written as $h(p, q)$. Why h is chosen rather than H will become apparent later:

$$\mathcal{N}_\nu \int \exp\{i \int [p\dot{q} - \dot{G}(p, q) - h(p, q)] dt\} \exp\{-\frac{1}{2\nu} \int (d\sigma^2/dt^2) dt\} \mathcal{D}p \mathcal{D}q \quad (46)$$

Here $\nu \in \mathbb{R}^+$. This expression is still a formal path integral. A very important step on the way to a well-defined path integral is to realize that

$$\mathcal{N}_\nu \exp\{-\frac{1}{2\nu} \int (d\sigma^2/dt^2) dt\} \mathcal{D}p \mathcal{D}q =: d\mu_W^\nu \quad (47)$$

is a well-defined pinned Wiener measure. Mathematically one defines the Wiener measure $d\mu_W^\nu$ by setting $\int d\mu_W^\nu := [\exp\{\nu T \Delta_{LB}\}](p'', q'', q', p')$, where the last expression is the kernel of the exponentiated Laplace-Beltrami operator (or heat kernel).

As was pointed out in section (2.6) the Wiener measure is concentrated on Brownian motion-like paths. So some integrals like $\int q dp$ have to be understood as stochastic integrals, and the Stratonovich rule is adopted for them. Then

$$\int \exp\{-i \int [q dp + d\tilde{G}(p, q) + h(p, q) dt]\} d\mu_W^\nu \quad (48)$$

is a well-defined path integral and expression (46) can be given meaning by equating it to this path integral (with $\tilde{G} = -(G + pq)$).

The question is: What happens in the limit of diverging diffusion constant ν (where the regularization formally becomes unity)?

If this limit leads to a well-defined path integral then further questions follow: What kind of path integral is it, i.e. to which states is it connected?

¹⁹For multiply connected manifolds the phase factor carries the Aharonov-Bohm phase.

What is the relation of the classical and quantum Hamiltonian²⁰?

For the case of the flat and spherical metric these questions were answered by Daubechies/Klauder [39] and for the hyperbolic metric by Daubechies/Klauder/Paul (DKP) [26]:

In the limit of diverging diffusion constant ν the path integral (48) reduces to the coherent state matrix element $\langle p''q'' | \exp\{-iT\mathcal{H}\} | p'q' \rangle$ of the unitary time-evolution operator and the specific metric determines the coherent states in question. The flat metric is inevitably connected with the coherent states of the Heisenberg-Weyl group, and in the canonical, Cartesian form, it is connected to the canonical coherent states. The spherical metric is associated with the coherent states of the SU(2) group and the hyperbolic metric leads to the coherent states of the affine group. And with each group comes a set of quantum kinematical operators. Thus one can say that the choice of geometry augmenting the classical phase space manifold determines the quantum kinematical operators uniquely! Furthermore, the classical Hamiltonian that goes with the quantum Hamiltonian $\mathcal{H}$ is given by the lower symbol (explaining why h was used earlier instead of H to characterize the classical Hamiltonian).

For a better understanding of how these results emerge it is instructive to point out the idea of the proof which is similar in the three cases and shall be illustrated for the affine group since this group will be of the most interest in the later work.

The construction of the coherent state path integral with Wiener measure for the affine group proceeds in two steps. First, the path integral for a zero Hamiltonian is derived; second, a non-zero Hamiltonian is introduced. The key to the first task is a linear (complex) polarization condition, an idea borrowed from the program of “geometric quantization”. Why this polarization condition has to be linear will become apparent. Observe, that the minimum uncertainty states $|\eta\rangle$ from section (2.3) are extremal weight vectors which fulfill $(Q - 1 + i\beta^{-1}D)|\eta_\beta\rangle = 0$. Using the definition of the affine coherent states, this leads directly to a linear first order differential operator

$$B = -iq^{-1}\partial_p + 1 + \beta^{-1}q\partial_q \quad (49)$$

which annihilates the coherent state overlap $\langle pq|rs\rangle = 2^{-2\beta}(qs)^{-\beta}[(q^{-1} + s^{-1}) + i\beta^{-1}(p - r)]^{-2\beta}$. As a positive-definite function the overlap can be

²⁰Remember: Due to the formal nature of the standard path integrals there were ambiguities concerning this point. But a well-defined path integral has to select a specific connection between the classical and quantum Hamiltonian.

taken as a reproducing kernel and thus B annihilates the whole reproducing kernel Hilbert space $\mathcal{C}_\beta$ built from the overlap [27]. This condition $B\mathcal{C}_\beta = 0$ is the linear complex polarization condition referred to above.

At this point specialize to a parameter range $\beta > 1/2$ for the fiducial vector $|\eta\rangle$. Then the fiducial vector admissibility condition (26) is fulfilled and the representation is square-integrable. This means $\mathcal{C}_\beta$ is a subspace of $L^2(M_+)$. Define the operator

$$\begin{aligned} A &:= \frac{1}{2}\beta B^\dagger B \\ &= \frac{1}{2}\{-\beta^{-1}\partial_q q^2 \partial_q - \beta q^{-2}\partial_p^2 - 1 + \beta - 2i\beta q^{-1}\partial_p\} \end{aligned} \quad (50)$$

Naturally one has $A\mathcal{C}_\beta = 0$. DKP [26] showed that A is a non-negative, self-adjoint operator²¹. For $\beta > 1/2$ the operator A has a discrete eigenvalue 0, which is clear since A annihilates the subspace $\mathcal{C}_\beta$. Furthermore, the spectrum of A was derived by reducing the 2-dimensional problem to a 1-dimensional one involving the Morse operator, the spectrum of which is well-known [40]. Thus, DKP additionally showed that 0 is an isolated eigenvalue and concluded that the semigroup $\exp\{-\nu T A\}$ strongly converges to the projection operator P_0 onto the subspace $\mathcal{C}_\beta$ in the limit of diverging diffusion constant ν . This ensures convergence of the kernels in a distributional sense: $[\exp\{-\nu T A\}](p'', q'', p', q') = \exp\{-\nu T A\}\delta(p-p')\delta(q-q')|_{p=p'', q=q''} \rightarrow [P_0](p'', q'', p', q') = (1 - \frac{1}{2\beta})(2\pi)^{-1}\langle p'' q'' | p' q' \rangle$. It is the author's point of view that a discrete eigenvalue 0 is already enough and that the question about a gap separating 0 from the rest of the spectrum is not relevant. Assume that there is no gap, i.e. the continuous spectrum starts at 0, too. Then, ignoring degeneracy, $\exp\{-\nu T A\} = \sum \exp\{-\nu T n\}|n\rangle\langle n| + \int \exp\{-\nu T \lambda\}|\lambda\rangle\langle\lambda|d\lambda$ and in the limit $\nu \rightarrow \infty$ this reduces to $|0\rangle\langle 0| = P_0$ since the single non-zero point in the integral, coming from the continuous "eigenvalue" 0, does not affect the zero result of the integral. As a next step, pointwise convergence of the kernels was established. Finally the Feynman-Kac-Stratonovich representation of the operator $\exp\{-\nu T A\}$ (or its kernel,

²¹To compare the work of DKP to the one here it must be noticed that they generally chose the a - b notation rather than the p - q notation. At the point where they talk about the connection between their a - b notation and a certain p - q notation one has to interchange p and q again to match the notation in the present work. Another difference is their choice of the one-parameter family $|\eta_\beta\rangle$. Whereas they chose to set the mean of Q equal to β the choice is here $Q=1$ and so their β is physically different from the one here but mathematically things turn out to be the same apart from some factors β here and there. See appendix (A.1).

to be more precise) was derived²². It is

$$\mathcal{N}_\nu \int \exp\{i \int [p\dot{q}] dt\} \exp\{-\frac{1}{2\nu} \int (\beta q^{-2} \dot{q}^2 + \beta^{-1} q^2 \dot{p}^2) dt\} \mathcal{D}p \mathcal{D}q \quad (51)$$

which is Eq. (46) for the special case of the hyperbolic metric and zero Hamiltonian. Equation (51) can be turned into a well-defined path integral by introducing the Wiener measure

$$d\mu_W^\nu := \mathcal{N}_\nu \exp\{-\frac{1}{2\nu} \int (\beta q^{-2} \dot{q}^2 + \beta^{-1} q^2 \dot{p}^2) dt\} \mathcal{D}p \mathcal{D}q \quad (52)$$

Combining this with the foregoing arguments, it was proved that the path integral for zero Hamiltonian

$$\int \exp\{-i \int q dp\} d\mu_W^\nu \quad (53)$$

is proportional to the coherent state overlap in the limit of diverging diffusion constant. At this point the requirement of a linear (complex) polarization condition can be explained: Only a linear polarization condition for the fiducial vectors leads to a first-order differential operator B and, ergo, to a second-order differential operator A . The Wiener measure emerges only because A is second-order. A higher-order polarization condition for the fiducial vector would not lead to a Wiener measure path integral.

Finally, moving along the same lines outlined for zero Hamiltonian, DKP showed the well-definedness of the coherent state path integral with Wiener measure for a wide (dense) class of Hamiltonians (at least polynomial). This proof is not so important for the work at hand and the reader is referred to [39] [26]. Suffice it to say, the classical Hamiltonian associated with the quantum Hamiltonian turns out to be determined by the lower symbol.

The mathematical side of the new coherent state path integral with Wiener measure has been discussed for the three 2-dimensional homogeneous spaces of constant curvature and the well-definedness of the integral has been established. But the path integral

²²The calculation is lengthy, see subsection (3.2.3) where it is done for a modified operator A_ε . Set $\varepsilon = 0$ to get back to the operator A

$$\begin{aligned}
& \langle p'' q'' | \exp\{-iT\mathcal{H}\} | p' q' \rangle \\
&= 2\pi(1 - \frac{1}{2\beta})^{-1} \lim_{\nu \rightarrow \infty} \int \exp\{-i \int [q dp + dG(p, q) + h(p, q) dt]\} d\mu_W^\nu \\
&=: 2\pi(1 - \frac{1}{2\beta})^{-1} \lim_{\nu \rightarrow \infty} \mathcal{N}_\nu \int \exp\{-i \int [q \dot{p} + \dot{G}(p, q) + h(p, q)] dt\} \\
&\quad \times \exp\{-\frac{1}{2\nu} \int (d\sigma^2/dt^2) dt\} \mathcal{D}p \mathcal{D}q \tag{54}
\end{aligned}$$

has more to offer.

First, observe that the metric in the regularization factor²³ gives physical meaning to the coordinatized expression for the action $-\int q dp + dG + h(p, q) dt$, which otherwise is simply not there. (Compare the discussion in section (2.6) where the concept of a shadow metric on the classical phase space was introduced.) In the standard path integrals this question did not have to be raised since the lattice, on which the formal integral was defined, automatically interpreted the Hamiltonian.

Second, by virtue of the Stratonovich interpretation of the stochastic integrals, canonical (coordinate) transformations can be performed. The coherent states are (apart from relabeling and unimportant phase factors) unaffected by these transformations [39]. The quantization is truly geometric in nature. (Since the term “geometric quantization” has already been claimed, it is for obvious reasons sometimes called “metrical quantization”). The connection of a phase space manifold with a specific shadow metric and the groups generating the coherent states are fundamental and independent of the coordinatization. The only thing that changes under canonical (coordinate) transformations is the way the groups are parameterized. Generally, the group will not be in group coordinates of the first or second kind anymore).

Finally note that a generalization of the above to 2-dimensional phase space manifolds with arbitrary choice of geometry is possible [41][27]. Since it will not come up again in later sections only the result is given here. The path integral is

²³or taking the more mathematical approach: the metric that enters the definition of the Wiener measure through the Laplace-Beltrami operator

$$\begin{aligned}
& \langle \xi'', T | \xi', 0 \rangle = \langle \xi'' | e^{-i\mathcal{H}T} | \xi' \rangle \\
& = \lim_{\nu \rightarrow \infty} \mathcal{N}_\nu \int \exp\{i \int [a_j(\xi) \dot{\xi}^j - h(\xi)] dt\} \exp\{-\frac{1}{2\nu} \int g_{jk}(\xi) \dot{\xi}^j \dot{\xi}^k dt\} \\
& \quad \times \exp\{\frac{\nu}{4} \int \sqrt{g(\xi)} \varepsilon^{jk} f_{jk}(\xi) dt\} \prod_t \sqrt{g(\xi)} d\xi^1 d\xi^2
\end{aligned}$$

where ξ^j , $j = 1, 2$ are coordinates, g_{jk} the metric tensor, g its determinant, a_j is a vector, $f_{jk} = \partial_j a_k - \partial_k a_j$ the field tensor and h the Hamiltonian.

In general there will be no group to define the coherent states. The metric tensor g_{jk} added to phase space may well differ from the metric induced by the coherent states (whereas in the case of a homogeneous space they coincided). The symplectic form need no longer be proportional to the volume element. Still, the states $|\xi\rangle$ are coherent states (so $\mathbb{1} = \int |\xi\rangle \langle \xi| \sqrt{g(\xi)} d\xi^1 d\xi^2$, resolution of unity) and the quantum Hamiltonian is associated to the classical Hamiltonian given by the lower symbol: $\mathcal{H} = \int h(\xi) |\xi\rangle \langle \xi| \sqrt{g(\xi)} d\xi^1 d\xi^2$.

3 Weak Coherent States and the Weak Coherent State Path Integral

If the second defining property of coherent states, the resolution of unity, is relaxed in a certain way one obtains what will be called the Klauder states, an even further generalization of the coherent states. Those Klauder states which are not coherent states in the old sense will be named weak coherent states. The definition of Klauder states and, therefore, weak coherent states will be given in section (3.1). The main part of this work is the study of the existence of path integrals for weak coherent states which follows in section (3.2).

Before this program is carried out some motivating remarks are in place about why a generalization of coherent states and their path integrals are of interest.

Affine quantum gravity

The spatial part of the space-time metric $g_{\mu\nu}(x)$ is strictly positive definite. Upon quantization this 3×3 matrix-valued field variable becomes a field operator which preserves this property and, hence, can not be an unbounded operator satisfying canonical commutation relations. The positivity property strongly suggests affine commutation relations, the simple analog of which is Eq. (21) for one degree of freedom. Whereas in this simple case affine commutation relations follow from the canonical commutation relations by multiplication with Q , this is not true in the case of fields (due to a certain infinite product and infinite rescaling). To choose affine commutation relations means then to discard canonical ones and the quantization is noncanonical [42][43].

Taking the affine field operators as generators one can define coherent states, or to be more cautious, Klauder states, since effectively it turns out that the overlap will in the simplest case of constant fields have a functional form which is analogous to the one for the affine weak coherent states. In this way one is naturally led to study the problem for the affine weak coherent states. Whereas in the latter case one can explicitly describe the family of fiducial vectors ($|\eta_\beta\rangle$) this is not possible for the general case of non-constant fields. There the overlap is central to the analysis [42] and the fiducial vector is implicitly contained in the functional form of the overlap.

The natural question is whether there is a path integral representation

for this overlap. For the special case of constant fields, which is equivalent to the affine (weak) coherent state problem for one degree of freedom, it is known that there is a well-defined path integral for a certain parameter range ($\beta < 1/2$), namely the one presented in section (2.7). On the other hand it turns out that for the parameter range $0 < \beta \leq 1/2$ the states do not admit a resolution of unity and thus they are weak coherent states.

Hence, an important step on the way to establish a path integral representation for the case of the affine fields is to prove the existence of a well-defined affine weak coherent state path integral for one degree of freedom. Certainly there can be no conventional way to define this path integral since the construction of the standard path integral requires a resolution of unity. But the concepts of the coherent state path integral with Wiener measure might be the key for an approach to a weak coherent state path integral. This was proposed by Klauder [27][42] and it is the aim of this work to shed some light on the existence of weak coherent state path integrals.

3.1 Weak coherent states

If one relaxes the second property which defines coherent states, namely that they possess a resolution of unity, and postulates only that the new states span the Hilbert space one arrives at a generalization called Klauder states. In mathematical terms Klauder states are defined by

1) Continuity: The states $|l\rangle$ are a strongly continuous vector-valued function of the label l .

2) Completeness: The family of vectors $(|l\rangle)$ is total, i.e. the closed linear span of $(|l\rangle)$ is the whole Hilbert space $\mathfrak{H}$.

The Klauder states which are not coherent states, i.e. which do not have a resolution of unity, are named weak coherent states.

The first question is: Is the class of Klauder states really bigger than the class of coherent states? Intuitively this seems clear, but it is proved once one can specify one set of weak coherent states explicitly. This is done subsequently and the weak coherent states in question will be group-defined.

From the discussion in section (2.3) it is clear that groups for which left- and right-invariant group measure coincide can not create weak coherent states, since all their representations are square integrable. Of the three group-defined sets of coherent states - canonical, spin and affine - which were given a closer look at (and which are associated with a flat, spherical

and hyperbolic phase space respectively in the path integral) only the affine ones can possibly be weak coherent states. Equation (26) states a fiducial vector admissibility condition which must be fulfilled to ensure a resolution of unity for affine coherent states, namely the parameter β must be bigger than $1/2$ ²⁴.

Choosing a fiducial vector from the set of non-admissible vectors [Eq. (23)], i.e.

$$\eta_\beta(x) = N_\beta x^{\beta-1/2} e^{-\beta x} \quad (55)$$

for $\beta \leq 1/2$, one indeed obtains weak coherent states. Joint continuity in the overlap [Eq. (25)]

$$\langle pq|rs\rangle = (qs)^{-\beta} 2^{-2\beta} [(q^{-1} + s^{-1}) + i\beta^{-1}(p-r)]^{-2\beta} \quad (56)$$

is clear. Also, the states span the whole space which follows from the irreducibility of the group representation.

No studies of other weak coherent states have been made so far, but it seems highly probable that every group with similar properties can define weak coherent states when the fiducial vector is chosen such that the representation is non-square integrable. But whether they or possibly non-group-defined weak coherent states are of physical interest has yet to be determined.

3.2 The affine weak coherent state path integral

In section (2.7) the coherent state path integral with Wiener measure was derived. For the affine coherent states the parameter range for the fiducial vectors $|\eta\rangle$ had to be limited to $\beta > 1/2$. This ensured the resolution of unity and the discrete (isolated) eigenvalue 0 for the operator A needed in the course of the development of the path integral.

Things change very much when the parameter range is $0 < \beta \leq 1/2$. As was shown in section (3.1) the states generated by the affine group acting on fiducial vectors with $\beta \in (0, 1/2]$ are no longer coherent states, but weak coherent states.

It is clear that a weak coherent state path integral - should it exist - can not be constructed in the usual way, since the standard procedure of time slicing requires a resolution of unity. One way is to start from the

²⁴With the connection to the Lobachevsky half-plane as the underlying phase space manifold this amounts to the requirement that the (scalar) curvature of the hyperbolic manifold, which is $R = -2/\beta$, must be less than -4

well-defined path integral with Wiener measure, and to try to adapt it to the new situation of weak coherent states.

For $0 < \beta \leq 1/2$ the coherent state overlap is no longer in $L^2(M_+)$. Consequently, the reproducing kernel Hilbert space $\mathcal{C}_\beta$ is not in the Hilbert space $L^2(M_+)$, and so 0 can not be a discrete eigenvalue any more. In fact it will be shown that only for $\beta = 1/2$ the vectors in $\mathcal{C}_\beta$ are generalized eigenvectors to the eigenvalue 0. The question why 0 is not in the (continuous) spectrum in spite of the fact that still $A\mathcal{C}_\beta = 0$ for all $\beta \in (0, 1/2)$ will be investigated.

If 0 is in the continuous spectrum of a non-negative operator X one has to face the problem that $\lim_{\nu \rightarrow \infty} e^{-\nu TX} = 0$ [$e^{-\nu TX} = \int_0^\infty e^{-\nu T\lambda} |\lambda\rangle\langle\lambda| d\lambda \xrightarrow{\nu \rightarrow \infty} 0$ since only one point in the integrand “survives” in the limit (namely $\lambda = 0$) which does not change the value of the integral]. Thus, it is not possible to arrive at a valid path integral representation with Wiener measure in the same way as before since the reduction to a projection operator onto the ground state in the limit of diverging diffusion constant was crucial in its construction [section (2.7)].

Klauder [27] proposed a procedure to prevent this collapse onto 0 by rescaling with a ν -dependent factor before taking the limit (which amounts to putting a δ -function weight on the “eigenvalue” 0). He assumed that this procedure to isolate the space $\mathcal{C}_\beta$ could be carried out for the whole parameter range $0 < \beta \leq 1/2$. Since $0 \in \text{spec}(A)$ only for $\beta = 1/2$ this idea, which will henceforth be called the spectral approach, can and will be developed in this case exclusively.

In the following subsection (3.2.1) the isolating procedure is first explained on a toy model. The procedure is then developed for an arbitrary self-adjoint operator. After the questions about the spectrum of the operator A have been clarified, the general formulae are applied to the affine case for $\beta = 1/2$. The corresponding weak coherent state path integral for zero Hamiltonian is formulated. Dynamics and symbols are introduced in subsection (3.2.2). The case $0 < \beta < 1/2$ is treated afterwards with a different method, namely a regularization approach. Again a zero Hamiltonian is assumed first in subsection (3.2.3) before dynamics are introduced in subsection (3.2.4).

3.2.1 Spectral approach

To explain the procedure to isolate the space $\mathcal{C}_\beta$ in the spectral approach a simple example, given by Klauder [27], is good for illustration:

Toy example

Let $\tilde{A} := \frac{1}{2}\tilde{B}^2$, $\tilde{B} := -i\partial_x$ on $L^2(\mathbb{R})$. Then $\tilde{B}$ is self-adjoint, and so is $\tilde{A}$ with $\tilde{A} \geq 0$. The eigenvalue $\lambda = 0$ lies in the continuous spectrum of $\tilde{A}$. The aim is to isolate the space of generalized eigenfunctions (non-square integrable functions) belonging to the eigenvalue $\lambda = 0$ of the operator $e^{-\nu T \tilde{A}}$ in the limit $\nu \rightarrow \infty$.

Klauder says at this point: “The desired space of functions is composed of those for which $\psi(x) = \text{const}$, namely a one-dimensional space. We can choose the reproducing kernel for this space to be identically one.”

A few comments on this statement are helpful, since one can easily find a two dimensional space of solutions of $\tilde{A}\psi = 0$, the space spanned by $\{1, x\}$, and ask the question: Why does the second solution x not come into the game?

The answer to this is twofold. First, one has to understand what is meant by “the desired space of functions”. Why can one simply talk about such a space and ignore the other solution? Since the goal is to find a valid path integral representation for only this kernel one should almost certainly be justified in ignoring the other solution. The kernel (1 in this case) is the starting point! To make this clear consider the original problem. Whatever the (weak coherent state) path integral may be, the coherent state overlap must be the kernel for zero Hamiltonian ($\langle pq|e^{-iHt}|p'q'\rangle = \langle pq|p'q'\rangle$ for $H = 0$ independent of what the path integral looks like). So what one is trying to do is to “inflate” the simple coherent state overlap to a meaningful path integral. The same is valid for the toy example. The question is how to build a well-defined path integral representation of 1, but then it is unlikely that an “ x -part” should appear in the process.

If such an “ x -part” really appeared it would have to cancel exactly or vanish when taking limits. So one can still insist on the question about a second solution. There is even more reason to do so, since later things will be approached from the other direction, i.e. some path integral will be taken as a starting point and its reduction (in the limit of diverging diffusion constant) to the kernel will be shown.

But indeed, the “second solution” can not appear and there is quite a subtle point to be learned on the way.

Before tackling the main problem remember that often some solutions of certain equations (differential or other) are ruled out by additional requirements like positivity, square integrability etc. So one could ask the new question: What could be such an additional condition here to prevent the

“second solution” from entering the picture?

This issue is addressed first on Hilbert space. Assume $B^\dagger B\psi = 0$. Take $0 = \langle \psi | B^\dagger B\psi \rangle = \langle B\psi | B\psi \rangle$, so $B\psi = 0$ and thus $B^\dagger B\psi = 0 \Leftrightarrow B\psi = 0$ and in this case no more solutions can exist.

For the toy example, change the kernel 1 to an (unnormalized) Hilbert space vector by multiplying it with $e^{-\frac{1}{2}\varepsilon x^2}$ (the limit $\varepsilon \rightarrow 0$ will be taken as the last step). Asking the question which operator annihilates this new kernel would lead to an ε -modified version of $\tilde{B}$ and, therefore, of $\tilde{A}$. (And, indeed, this is the right way to proceed. The kernel is the starting point and determines $\tilde{B}_\varepsilon$ and $\tilde{A}_\varepsilon$!) One gets $\tilde{B}_\varepsilon = -i\partial_x - i\varepsilon x$ and $\tilde{A}_\varepsilon = \frac{1}{2}(-\partial_x^2 + \varepsilon^2 x^2 - \varepsilon)$. By the little Hilbert space theorem above there can not be more solutions than $e^{-\frac{1}{2}\varepsilon x^2}$ and the limit will certainly take things back to the original kernel 1.

The regularizing approach in subsection (3.2.3), which will solve the problem of the affine weak coherent states for the parameter range $0 < \beta < 1/2$, is just such an ε -modification. In whatever way the modification might complicate things, one certainly does not have to worry about a “second solution”. But there is still more to say:

What would this procedure look like for the “second solution” x ? Consider the unnormalized Hilbert space vector $x e^{-1/2\varepsilon x^2}$. The ε -modified version of $\tilde{B}$ would be $\tilde{B}_\varepsilon = -i\partial_x - i\varepsilon x + i/x$. But, this would not reduce to $\tilde{B}$ in the limit $\varepsilon \rightarrow 0$, or else the second and first solution would be the same. For $\tilde{B}_\varepsilon$ to make sense one has to impose $\psi(0) = 0$ for the solutions! This will be called a Dirichlet boundary condition (D.B.C.), although 0 is, strictly speaking, not a boundary here. In the limit $\tilde{A}_\varepsilon = \frac{1}{2}\tilde{B}_\varepsilon^\dagger \tilde{B}_\varepsilon = \frac{1}{2}(-\partial_x^2|_{D.B.C.} + \varepsilon^2 x^2 - 3\varepsilon)$ does not reduce to $\tilde{A}$ since the ε -modified version carries the D.B.C. and is positive definite only because the D.B.C. excludes the ground state of this oscillator (see e.g. [34]).

So the final answer is that by asking the naive question about a “second solution” the additional constraint of boundary conditions was overlooked. Since a path integral is a solution to a differential equation, it implicitly carries a boundary condition. That is why neither in the toy example nor in the case of the affine weak coherent states one will have to worry about more solutions than the ones already given by $\tilde{B}\psi = 0$ (or later $B\psi = 0$). A solution of $B^\dagger B\psi = 0$ for which $B\psi \neq 0$ would carry a different boundary condition.

This interesting point is connected to the problem of operator extensions (Friedrichs’ extension, Krein extension etc. [44]).

Returning to the toy example one finds by direct calculation²⁵ (or with the aid of the Mehler formula [29] for the oscillator path integral followed by taking the oscillator frequency to 0):

$$\langle x'' | e^{-\nu T \tilde{A}} | x' \rangle = e^{-\nu T \tilde{A}(x)} \delta(x - x') \big|_{x=x''} = \frac{1}{\sqrt{2\pi\nu T}} e^{-\frac{(x''-x')^2}{2\nu T}} = \int d\mu_W^\nu(x)$$

It is the factor $(2\pi\nu T)^{-1/2}$ which is responsible for the collapse onto 0. By the rescaling with $\sqrt{2\pi\nu T}$, one gets

$$\lim_{\nu \rightarrow \infty} \sqrt{2\pi\nu T} \int d\mu_W^\nu(x) = \lim_{\nu \rightarrow \infty} e^{-\frac{(x''-x')^2}{2\nu T}} = 1 \quad (57)$$

which gives the correct reproducing kernel for the desired subspace.

Also, notice that

$$e^{-\frac{(x''-x')^2}{2\nu T}} = \frac{\int d\mu_W^\nu(x)}{\int_{x'=0}^{x''=0} d\mu_W^\nu(x)} \quad (58)$$

is a self-consistent way to determine the proper rescaling factor.

The general case

Let X be a non-negative, self-adjoint operator on a certain Hilbert space and assume 0 is in its continuous, but not in its discrete spectrum. The operator X generates a semigroup $e^{-\nu T X}$ which has a spectral representation $e^{-\nu T X} = \int_0^\infty e^{-\nu \lambda T} d\mathbb{E}(\lambda)$ or $\langle x'' | e^{-\nu T X} | x' \rangle = \int_0^\infty e^{-\nu \lambda T} d\langle x'' | \mathbb{E}(\lambda) | x' \rangle$.

Since only rather well-behaved potentials will eventually be of interest, the reasonable assumption is made that the measure $d\langle x'' | \mathbb{E}(\lambda) | x' \rangle$ has an absolutely continuous, but no singularly continuous part. Then the spectral family can be written as a (weighted) integral over one-dimensional projection operators $\mathbb{E}(\lambda) = \int_{-\infty}^\lambda |E\rangle \langle E| \rho(E) dE$ ²⁶. If the generalized eigenstates $|E\rangle$ are δ -orthonormalized, then $\rho(E) = 1$. This follows from the fact that every operator $\mathbb{E}(\lambda)$ is a projection operator, and hence $[\mathbb{E}(\lambda)]^2 = \int_{-\infty}^\lambda \int_{-\infty}^\lambda |E\rangle \langle E| E' \rangle \langle E' | \rho(E) \rho(E') dE dE' = \int_{-\infty}^\lambda \int_{-\infty}^\infty |E\rangle \delta(E - E') \langle E' | \rho(E) \rho(E') dE dE' = \int_{-\infty}^\lambda |E\rangle \langle E| \rho^2(E) dE = \mathbb{E}(\lambda) = \int_{-\infty}^\lambda |E\rangle \langle E| \rho(E) dE$. Since

²⁵Fourier representation of the δ -function and action of the operator will lead to simple Gaussian integrals.

²⁶For a singularly continuous measure this would not be possible: $\mu_{sc}(x) = \int_{-\infty}^x d\mu_{sc}(y) \neq \int_{-\infty}^x (d\mu_{sc}/dy) dy = 0$ since $d\mu_{sc}/dy = 0$ almost everywhere.

ρ is a non-negative function (and not identically zero almost everywhere) $\rho(E) \equiv 1$ follows.

The matrix element of $e^{-\nu TX}$ can then be written as

$$\langle x'' | e^{-\nu TX} | x' \rangle = \int_0^\infty e^{-\nu \lambda T} \psi_\lambda(x'') \psi_\lambda^*(x') \rho(\lambda) d\lambda \quad (59)$$

and the ψ_λ are continuous in λ . Moreover, ρ - being part of the measure - is at least right-continuous and for δ -orthonormalized wavefunctions $\rho(\lambda) \equiv 1$.

The goal is to find the rescaling factor which saves equation (59) from becoming trivial in the limit of diverging diffusion constant ν . Since for very large ν the factor $e^{-\nu \lambda T}$ suppresses everything but very small λ , the behavior of $f_{x',x''}(\lambda) := \psi_\lambda(x'') \psi_\lambda^*(x') \rho(\lambda)$ is all that matters. To give an example assume that $f_{x',x''}(\lambda) \propto \lambda^a$ for small λ . Then the proper rescaling factor can be determined such that $\int_0^\infty e^{-\nu \lambda T} \psi_\lambda(x'') \psi_\lambda^*(x') \rho(\lambda) d\lambda$ will not collapse to 0 but instead go to the desired $\psi_0(x'') \psi_0^*(x')$. In the example this factor is

$$\int_0^\infty d\lambda \lambda^a e^{-\nu \lambda T} = \frac{\Gamma(a+1)}{(\nu T)^{a+1}} \quad (60)$$

which goes to 0 for $\nu \rightarrow \infty$. After rescaling with the inverse one gets $\frac{(\nu T)^{a+1}}{\Gamma(a+1)} \lambda^a e^{-\nu \lambda T} \xrightarrow{\nu \rightarrow \infty} \delta(\lambda)$, which is the desired δ -function weight on 0.

As in the toy example, the rescaling factor can be computed self-consistently and the general formula becomes

$$\lim_{\nu \rightarrow \infty} \frac{\int_0^\infty e^{-\nu \lambda T} \psi_\lambda(x'') \psi_\lambda^*(x') \rho(\lambda) d\lambda}{\int_0^\infty e^{-\nu \lambda T} \psi_\lambda(0) \psi_\lambda^*(0) \rho(\lambda) d\lambda} \xrightarrow{\nu \rightarrow \infty} \psi_0(x'') \psi_0^*(x') \quad (61)$$

The last expression, $\psi_0(x'') \psi_0^*(x')$, is the kernel of the desired projection operator onto the ground state. The convergence is in a distributional sense (denoted by the symbol $\rightarrow$). If the functional form of $\psi_0(x'') \psi_0^*(x')$ is known to be continuous then the convergence is pointwise.

Observe in the example with $f_{x',x''}(\lambda) = \lambda^a$ one must have $a > -1$ or else the rescaling factor would be identically 0 (since the integral would be infinity). But since the rescaling factor can be determined self-consistently, i.e. by the denominator of equation (61), which always exists, there is no hidden “trap” to look out for. Moreover the evaluation of the denominator need not necessarily be at the point $x'' = x' = 0$. It could be at any point $x'' = x' = b$, $b \in \mathbb{R}$ or even $b = \pm\infty$ as long as the function $\psi_\lambda(x)$ is not 0 at b . Whatever gives the easiest result is the preferred choice. Sometimes this choice even makes no difference as in the toy example above, since

the function there depends on $x'' - x'$ only. But in general there is an arbitrariness, but this arbitrariness exists for a good reason. Assume that some expression K serves as a reproducing kernel. Let a be a constant, then aK is just as good a reproducing kernel, but the inner product in the reproducing kernel Hilbert space must be redefined. Sometimes, on the other hand, there clearly is a preferred choice. Take the case of the affine coherent states overlap ($\beta > 1/2$). The overlap is a kernel for a reproducing kernel Hilbert space $\mathcal{C}_\beta$, but since there is a resolution of unity which gives rise to an integral definition of the inner product, the factor in front of the reproducing kernel is fixed (or at least it would not make much sense to define another inner product on $\mathcal{C}_\beta$). This factor used to be $(1 - \frac{1}{2\beta})(2\pi)^{-1}$ and appeared frequently in the affine coherent state path integral. For the case of the weak coherent state path integral ($\beta \leq 1/2$) on the other hand, the factor is no longer fixed, since the resolution of unity does not exist anymore.

The function $f_{x',x''}(\lambda) = \langle x'' | \delta(\lambda - X) | x' \rangle$ is connected to the Green's function and its determination proceeds as follows.

Define $\delta_+(y) := \lim_{\varepsilon \rightarrow 0} \int_0^\infty \frac{1}{2\pi} e^{iyt - \varepsilon t} dt = \lim_{\varepsilon \rightarrow 0} \frac{1}{2\pi} \frac{1}{\varepsilon - iy}$. Write $\frac{1}{\varepsilon - iy} = \frac{1}{\varepsilon - iy} \frac{\varepsilon + iy}{\varepsilon + iy} = \frac{iy}{\varepsilon^2 + y^2} + \frac{\varepsilon}{\varepsilon^2 + y^2}$ and since $\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^\infty \frac{\varepsilon dy}{\varepsilon^2 + y^2} = \int_{-\infty}^\infty \frac{du}{1+u^2} = \pi$ one has $\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon - iy} = i\mathcal{P}(\frac{1}{y}) + \pi\delta(y)$ where the principal value $\mathcal{P}$ is defined by $\int \mathcal{P}[\frac{f(y)}{y}] dy := \lim_{\varepsilon \rightarrow 0} (\int_\varepsilon^\infty + \int_{-\infty}^{-\varepsilon}) \frac{f(y)}{y} dy$. Hence, the δ_+ -function can be expressed as $\delta_+(y) = \frac{1}{2}\delta(y) + \frac{i}{2\pi}\mathcal{P}(\frac{1}{y})$.

The Green's function $G = \lim_{\varepsilon \rightarrow 0} \int_0^\infty e^{i(E+i\varepsilon)T} \langle x'' | e^{-iHT} | x' \rangle dT$ is, after integration, seen to be $G = \lim_{\varepsilon \rightarrow 0} \langle x'' | \frac{i}{(E-H)+i\varepsilon} | x' \rangle$. By the above $\lim_{\varepsilon \rightarrow 0} \frac{i}{(E-H)+i\varepsilon} = \pi\delta(E-H) + i\mathcal{P}(\frac{1}{E-H})$ and so by taking the real part one has

$$\begin{aligned} \Re \int_0^\infty dT e^{i(E+i\varepsilon)T} \langle x'' | e^{-iHT} | x' \rangle &= \Re \langle x'' | \int_0^\infty dT e^{iT(E-H)-\varepsilon T} | x' \rangle \\ &= \Re \langle x'' | \frac{i}{(E-H)+i\varepsilon} | x' \rangle \xrightarrow{\varepsilon \rightarrow 0} \pi \langle x'' | \delta(E-H) | x' \rangle = \pi f_{x',x''}(E) \end{aligned} \quad (62)$$

If the function $f_{x',x''}(E)$ is known then the rescaling factor can be determined - at least in principle - with the aid of the denominator of Eq. (61).

The spectral approach works for all operators which possess the same properties as the operator X .

The affine case

The foregoing is now applied to the case of the affine weak coherent states [see especially sections (2.3), (2.7) and (3.1)].

First, it is necessary, to see in which situation the spectral approach can possibly work. DKP [26] gave the spectrum of the operator A [which was defined in Eq. (50)] as

$$\begin{aligned} \text{spec}(A) = & \{(\beta - \tfrac{1}{2})^2 - (\beta - \tfrac{1}{2} - n)^2; n \in \mathbb{N}, n < \beta - \tfrac{1}{2}\} \\ & \cup [(\beta - \tfrac{1}{2})^2, \infty) \end{aligned} \quad (63)$$

Thus, there is no discrete spectrum for $0 < \beta \leq 1/2$. And only for $\beta = 1/2$ does the continuous spectrum include 0.

The question is: Why is 0 not in the spectrum even if all functions $\psi \in \mathcal{C}_\beta$ fulfill $A\psi = 0$ for the whole parameter range $\beta > 0$?

An equation $A\psi = \alpha\psi$ need not imply $\alpha \in \text{spec}(A)$. If ψ is square integrable then α is in the discrete spectrum. But if ψ is not square integrable, then α may or may not be in the continuous spectrum. The criterium is the following: There must be an (unnormalized) square integrable sequence $(\tilde{\psi}_n) \in L^2(M_+)^{\mathbb{N}}$ for which $\tilde{\psi}_n(x) \rightarrow \psi(x)$ for $n \rightarrow \infty$ (at least in a distributional sense) such that $\psi_n := N_n \tilde{\psi}_n$ is a normalized sequence (and N_n is a normalization constant) and $\|(A - \alpha)\psi_n\| \rightarrow 0$.

A toy example is again helpful to see this before the case on the Lobachevsky plane is discussed.

Let $\psi_\alpha = e^{i\alpha x}$ where $\alpha = p + ip'$. Take $P = -i\partial_x$, then $P\psi_\alpha = \alpha\psi_\alpha$. The claim is nonetheless that $\alpha \notin \text{spec}(A)$, except when $p' = 0$.

Consider a special sequence first: $\psi_n = N_n e^{i\alpha x} e^{-x^2/n^2}$.

Calculation shows: $1 \stackrel{!}{=} \|\psi_n\|^2 = N_n^* N_n e^{n^2(p')^2/2} (\frac{\pi}{2})^{1/2} n$ and with that one computes (partial integration and Gaussian integrals): $\|(P - \alpha)\psi_n\|^2 = 1/n^2 + (p')^2 = \|(P - p)\psi_n\|^2 + \|p'\psi_n\|^2$ which does not go to 0 for $n \rightarrow \infty$ except for $p' = 0$. This proves that for $\alpha = p$, i.e. $p' = 0$, $\alpha \in \text{spec}(A)$.

To complete the proof that $\alpha \notin \text{spec}(A)$ for $p' \neq 0$ the above must be shown for every sequence $(\psi_n) \in L^2(M_+)^{\mathbb{N}}$ having the required properties. This will not be further studied here, but be discussed for the real problem with the operator A on the Lobachevsky half-plane.

It is clear now that, even if $A\psi(p, q) = 0$ for all $\psi \in \mathcal{C}_\beta$, one still has to find out whether there is a normalized sequence $(\psi_n :) \in L^2(M_+)^{\mathbb{N}}$ such that $\|A\psi_n\| \rightarrow 0$ for $n \rightarrow \infty$. Since $\psi(p, q) = \langle pq | (\sum_{j=1}^J \alpha_j |p_j q_j\rangle)$

(or limits of corresponding Cauchy sequences), it is enough to consider $\psi(p, q) = \langle pq | rs \rangle = (1/2)^{-2\beta} (qs)^{-\beta} [(q^{-1} + s^{-1}) + i\beta^{-1}(p - r)]^{-2\beta}$.

For large q this expression is proportional to $q^{-\beta}$, and since $0 < \beta \leq 1/2$, a regularization factor, which is effective at infinity, is required to produce Hilbert space vectors again. Now, for $0 < 4\beta \leq 1$: $\int_{-\infty}^{\infty} (c^2 + p^2)^{-2\beta} dp = \infty$ (where c is a constant), and one must in this case regularize in p as well, whereas for $1 < 4\beta < 2$ this extra regularization is not needed. A regularization in p makes a regularization in q for small q necessary. This leads, in turn, to more complicated terms.

The case $1 < 4\beta < 2$ will be examined first. Most of the details are transferred to the appendix, since they consist of lengthy calculations which offer no great insight. Likewise, the case $0 < 4\beta < 1$ is studied entirely in appendix (A.3). The case $\beta = 1/2$ will be treated separately, and since 0 will then turn out to be in the spectrum, the isolating procedure referred to earlier will be carried out.

Note: Since the interest lies in the n -dependence of every expression, constants are generally absorbed in a factor “*const*” which can change from line to line.

I) $1 < 4\beta < 2$

The goal is to show $0 \notin \text{spec}(A)$. The proof starts with a special sequence: $\psi_n(p, q) := N_n (1/2)^{-2\beta} (qs)^{-\beta} [(q^{-1} + s^{-1}) + i\beta^{-1}(p - r)]^{-2\beta} e^{-q/n}$, the norm of which is

$$\begin{aligned}
\|\psi_n\|^2 &= N_n^* N_n s^{-2\beta} 4^{2\beta} \int_0^\infty dq e^{-2q/n} q^{-2\beta} \\
&\quad \times \int_{-\infty}^\infty dp [(q^{-1} + s^{-1})^2 + \beta^{-2}(p - r)^2]^{-2\beta} \\
&= N_n^* N_n s^{-2\beta} 4^{2\beta} \int_0^\infty dq e^{-2q/n} q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \\
&\quad \times \int_{-\infty}^\infty dp [1 + \beta^{-2} p^2]^{-2\beta} \\
&= \text{const} \cdot N_n^* N_n \int_0^\infty dq e^{-2q/n} q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \\
&= \text{const} \cdot N_n^* N_n n^{1-2\beta} \int_0^\infty dq e^{-2q} q^{-2\beta} [(qn)^{-1} + s^{-1}]^{1-4\beta}
\end{aligned}$$

This expression will always be smaller than $\text{const} \cdot N_n^* N_n n^{1-2\beta}$, which is the

correct asymptotic behavior for large n . So $N_n^* N_n > n^{2\beta-1} \times \text{const}^{-1}$.

Note that in the step from line 1 to 2 the substitution $p \rightarrow p/(q^{-1} + s^{-1})$ was made. The resulting p -integral can be found in [45], page 297, integral #3.252, 11.

With $[+] := [\frac{1}{2}(q^{-1} + s^{-1}) + \frac{1}{2}i\beta^{-1}(p - r)]$ one finds in this case: $A\psi_n = \{-q^2 n^{-2} + 2\beta[+]^{-1} n^{-1} + (2 - 2\beta)qn^{-1}\}\psi_n$

Use $[-] := [+]^* = [\frac{1}{2}(q^{-1} + s^{-1}) - \frac{1}{2}i\beta^{-1}(p - r)]$ and $[] := [-][+] = \frac{1}{4}[(q^{-1} + s^{-1})^2 + \beta^{-2}(p - r)^2]$ and $[+] + [-] = (q^{-1} + s^{-1})$ then calculation of $\|A\psi_n\|^2$ yields:

$$\begin{aligned}
& \|A\psi_n\|^2 \\
&= N_n^* N_n \int_{-\infty}^{\infty} dp \int_0^{\infty} dq e^{-2q/n} q^{-2\beta} \left\{ \frac{1}{4}[(q^{-1} + s^{-1})^2 + \beta^{-2}(p - r)^2] \right\}^{-2\beta} \\
&\quad \times \{-q^2 n^{-2} + 2\beta n^{-1}[-]^{-1} + 2(1 - \beta)qn^{-1}\} \\
&\quad \times \{-q^2 n^{-2} + 2\beta n^{-1}[+]^{-1} + 2(1 - \beta)qn^{-1}\} \\
&= \int_0^{\infty} dq [-q^2 n^{-2} + 2(1 - \beta)qn^{-1}]^2 \int_{-\infty}^{\infty} dp \psi_n^* \psi_n \\
&\quad + \int_0^{\infty} dq \int_{-\infty}^{\infty} dp \{-2\beta q^2([+]^{-1} + [-]^{-1})n^{-3} + 4\beta^2[+][-]n^{-2} \\
&\quad + 4\beta(1 - \beta)q([+]^{-1} + [-]^{-1})n^{-2}\} \psi_n^* \psi_n \\
&= \int_0^{\infty} dq [-q^2 n^{-2} + 2(1 - \beta)qn^{-1}]^2 \int_{-\infty}^{\infty} dp \psi_n^* \psi_n \\
&\quad + \int_0^{\infty} dq \int_{-\infty}^{\infty} dp \{-2\beta q^2([+] + [-])[]^{-1} n^{-3} + 4\beta^2[] n^{-2} \\
&\quad + 4\beta(1 - \beta)q([+] + [-])[]^{-1} n^{-2}\} \psi_n^* \psi_n
\end{aligned}$$

One has to make sure that cancelations do not occur. The first summand (the “ q -part”) goes to a constant, whereas the second (the “[]-part”) goes to zero:

$$\begin{aligned}
& N_n^* N_n \int_0^\infty dq [-q^2 n^{-2} + 2(1-\beta)qn^{-1}]^2 \int_{-\infty}^\infty dp (N_n^* N_n)^{-1} \psi_n^* \psi_n \\
> n^{2\beta-1} \int_0^\infty dq [-q^2 n^{-2} + 2(1-\beta)qn^{-1}]^2 e^{-2q/n} q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \\
= \int_0^\infty dq [-q^2 + 2(1-\beta)q]^2 e^{-2q} q^{-2\beta} [(qn)^{-1} + s^{-1}]^{1-4\beta}
\end{aligned}$$

This will not go to zero as $n \rightarrow \infty$ but to a constant > 0 ! The second term is

$$\begin{aligned}
& N_n^* N_n \int_0^\infty dq e^{-2q/n} q^{-2\beta} \int_{-\infty}^\infty dp \{ -2\beta q^2 ([+] + [-]) []^{-1} n^{-3} \\
& \quad + 4\beta(1-\beta)q([+] + [-]) []^{-1} n^{-2} + 4\beta^2 [] n^{-2} \} []^{-2\beta} \\
= & \text{const} \cdot N_n^* N_n \int_0^\infty dq e^{-2q/n} q^{-2\beta} \\
& \quad \times \{ [-2\beta q^2 n^{-3} + 4\beta(1-\beta)qn^{-2}] (q^{-1} + s^{-1})^{-4\beta} \\
& \quad + 4\beta^2 (q^{-1} + s^{-1})^{-1-4\beta} n^{-2} \} \\
= & \text{const} \cdot N_n^* N_n n^{1-2\beta} \int_0^\infty dq e^{-2q} q^{-2\beta} \\
& \quad \times \{ [-2\beta q^2 n^{-1} + 4\beta(1-\beta)qn^{-1}] [(nq)^{-1} + s^{-1}]^{-4\beta} \\
& \quad + 4\beta^2 n^{-2} [(nq)^{-1} + s^{-1}]^{-1-4\beta} \}
\end{aligned}$$

and the bottom line will go to zero due to the $1/n$, $1/n^2$ terms.

It has been shown that convergence fails for this special sequence. As in the toy example above, this is not yet a complete proof. One has to show such behavior for all possible ψ_n which fulfill the necessary requirements. This can be found in appendix (A.3) where the case $0 < 4\beta < 1$ and the limiting case $4\beta = 1$ are treated as well.

The case $\beta=1/2$

In this case the claim is $0 \in \text{spec}(A)$ (so it is enough to find one special sequence ψ_n that will do the job):

$$\begin{aligned}
\|\psi_n\|^2 &= \text{const} \cdot N_n^* N_n \int_0^\infty dq e^{-2q} q^{-1} [(nq)^{-1} + s^{-1}]^{-1} \\
&= \text{const} \cdot N_n^* N_n \left\{ \int_0^c dq e^{-2q} q^{-1} [(nq)^{-1} + s^{-1}]^{-1} \right. \\
&\quad \left. + \int_c^\infty dq e^{-2q} q^{-1} [(nq)^{-1} + s^{-1}]^{-1} \right\} \\
&\leq \text{const} \cdot N_n^* N_n \left\{ \int_0^c dq q^{-1} (nq) + \text{const}(c) \right\} \\
&= \text{const} \cdot N_n^* N_n [nc + \text{const}(c)]
\end{aligned}$$

So $N_n^* N_n \geq \text{const} \cdot [nc + \text{const}(c)]^{-1}$ and this is the correct asymptotic behavior. Observe how this is not the limit of the usual $n^{2\beta-1}$ as $\beta \nearrow 1/2$, which would be constant. As q^{-1} is no longer integrable at $q = 0$, one picks up an extra term of $1/(nc + \text{const}(c)) \sim 1/n$ for large n .

It is alright not to bother about the “[]-part” in $\|A\psi_n\|^2$ which was going to 0 even in the case $0 < \beta < 1/2$ (compare above). Now, the “ q -part” will also go to 0 by virtue of the new $1/n$ -dependence:

$$\begin{aligned}
&N_n^* N_n \int_0^\infty dq [-q^2 n^{-2} + q n^{-1}]^2 e^{-2q/n} q^{-1} (q^{-1} + s^{-1})^{-1} \\
&= N_n^* N_n \int_0^\infty dq [-q + 1]^2 q e^{-2q} [(nq)^{-1} + s^{-1}]^{-1}
\end{aligned}$$

The integral goes to a constant again (trivially, since the “ q -factors” have taken care of the q^{-1}). But the whole expression goes to zero because of the $1/n$ -dependence of the normalization factors $N_n^* N_n$! This completes the proof that $0 \in \text{spec}(A)$ for $\beta = 1/2$.

Indeed, what was established is exactly what was expected from the knowledge of the spectrum of A given by DKP [Eq. (63)].

Whereas there is no way to isolate the space $\mathcal{C}_\beta$ with the spectral approach in the case $0 < \beta < 1/2$ (simply because 0 is not in the spectrum), this is possible for $\beta = 1/2$ since the operator A fulfills all the requirements for the application of the general theory developed earlier. This does not seem like a major advance compared to the DKP-article, which treated $\beta > 1/2$, but in the case $\beta = 1/2$ one is dealing with weak coherent states for the first time!

The spectral approach and the path integral for $\beta=1/2$

As the operator A possesses all required properties in the case $\beta = 1/2$ (non-negativity, self-adjointness, 0 is in the continuous, but not in the discrete spectrum), it follows from the general theory that the procedure to isolate the subspace $\mathcal{C}_{1/2}$ can be carried out. This is enough to ensure that a weak coherent state path integral can be constructed which will subsequently be done, at first for a zero Hamiltonian. Dynamics will be introduced in the following subsection.

For the the present problem a certain connection (established in [26]) between the operator A and the Morse operator H_{Morse} exists and makes the explicit functional form of the generalized eigenfunctions available. With the aid of the latter one can compute the rescaling factor explicitly. With these calculations at hand it turns out to be just a short step to study the limit of diverging diffusion constant explicitly as well, confirming the general theory.

The problem to find the eigenfunctions of the operator A is reduced to a problem on $L^2(\mathbb{R}^+)$ and then to a problem on $L^2(\mathbb{R})$ leading to the Morse operator:

$$\begin{aligned}
A\langle U(p, q)\phi|\psi\rangle &= \langle A^*U(p, q)\phi|\psi\rangle \\
&= \frac{1}{2}\langle [-\beta^{-1}\partial_q q^2\partial_q - \beta q^{-2}\partial_p^2 - 2i\beta q^{-1}\partial_p + \beta - 1]e^{ipQ}e^{-i\ln qD}\phi|\psi\rangle \\
&= \frac{1}{2}\beta^{-1}\langle e^{ipQ}e^{-i\ln qD}\{D^2 + iD + \beta^2Q^2 - 2\beta^2Q + \beta^2 - \beta\}(Q^{1/2}\phi')|\psi\rangle \\
&= \frac{1}{2}\beta^{-1}\langle e^{ipQ}e^{-i\ln qD}Q^{1/2}\{D^2 + \beta^2Q^2 - 2\beta^2Q + (\beta - \frac{1}{2})^2\}\phi'|\psi\rangle \quad (64)
\end{aligned}$$

Here $e^{i\ln qD}Qe^{-i\ln qD} = qQ$ and $[D, Q^\alpha] = -i\alpha Q^\alpha$ were used, so $DQ^{1/2} = Q^{1/2}(D - i/2)$ and $D^2Q^{1/2} = Q^{1/2}(D - i/2)^2$. Observe that in the first line the operator A on the Lobachevsky plane is not an abstract Hilbert space operator $\mathcal{A}$, but already a representation thereof. Thus, it acts merely as a linear functional on the inner product, and, because of anti-linearity in the first position, it comes inside complex conjugated.

Under the unitary transformation

$$(\tilde{U}\psi)(x) = e^{x/2}\psi(e^x) \quad (65)$$

the operator in braces in the last line of Eq. (64) [called H in DKP] is transformed to the Morse operator:

$$H_{Morse} = -\frac{d^2}{dx^2} + \beta^2(e^{2x} - 2e^x) + (\beta - \frac{1}{2})^2 \quad (66)$$

The Green's function associated with the Morse operator (and derived from a conventional configuration space path integral) is given by Grosche and Steiner ([30], 6.4.2, page 228).

$$\begin{aligned} & \int_0^\infty dT e^{i(E+i\varepsilon)T/\hbar} \int_{x(t')=x'}^{x(t'')=x''} \exp\left\{\frac{i}{\hbar} \int_{t'}^{t''} \left[\frac{m}{2} \dot{x}^2 - \frac{\hbar^2 V_0^2}{2m} (e^{2x} - 2\alpha e^x)\right] dt\right\} \mathcal{D}x \\ &= -\frac{im}{2\hbar V_0} \Gamma(1/2 + \sqrt{-2mE}/\hbar - \alpha V_0) \Gamma^{-1}(1 + 2\sqrt{-2mE}) e^{(x'+x'')/2} \\ & \quad \times W_{\alpha V_0, \sqrt{-2mE}/\hbar}(2V_0 e^{x'}) M_{\alpha V_0, \sqrt{-2mE}/\hbar}(2V_0 e^{x''}) \end{aligned}$$

Here M and W are Whittaker functions, the definition of which is:

$$\begin{aligned} M_{\mu,\nu}(z) &= e^{-z/2} z^{1/2+\nu} {}_1F_1(1/2 - \mu + \nu, 1 + 2\nu, z) \\ W_{\mu,\nu}(z) &= e^{-z/2} z^{1/2+\nu} U(1/2 - \mu + \nu, 1 + 2\nu, z) \end{aligned} \quad (67)$$

where ${}_1F_1$ and U are confluent hypergeometric functions (of the first and second kind):

$$\begin{aligned} {}_1F_1(a, b, z) &= \Gamma(b) \Gamma^{-1}(a) \Gamma^{-1}(b-a) \int_0^1 e^{zt} t^{a-1} (1-t)^{b-a-1} dt \\ U(a, b, z) &= \Gamma^{-1}(a) \int_0^\infty e^{-zt} t^{a-1} (1+t)^{b-a-1} dt \end{aligned}$$

In this work $\hbar = 1$, and furthermore choose $m = 1/2$, $\alpha = 1$, $V_0 = \beta$ and finally $\beta = 1/2$ ²⁷

Grosche/Steiner [30] and Grosche [46] show that the Green's function can also be written as²⁸:

$$\begin{aligned} G(x'', x', E) &= \frac{-i}{\pi^2} \int_0^\infty d\lambda \frac{\lambda \sinh 2\pi\lambda}{\lambda^2 - E} |\Gamma(i\lambda)|^2 e^{-(x'+x'')/2} \\ & \quad \times W_{1/2, i\lambda}(e^{x'}) W_{1/2, i\lambda}(e^{x''}) \end{aligned} \quad (68)$$

First replace $\int_0^\infty$ by $\frac{1}{2} \int_{-\infty}^\infty$. This is correct since the Whittaker functions have the property $W_{\mu,\nu} = W_{\mu,-\nu}$ and the rest of the integrand is even as well. If the integral is now regularized by the convention $E \rightarrow E + i\varepsilon$ one can go

²⁷This is a critical case; for further details consult Grosche[46], page 118.

²⁸The factor $-i$ is not present in [46] since Grosche defines the Fourier transformation with an imaginary factor; this rule is not being followed here.

to a contour integral from $-R$ to R on the real line and on a semi-circle over this (can be closed in the upper or lower half plane). By Cauchy's integral formula the contour integral is $2\pi i$ times the residue of the integrand (which has a simple pole). Unfortunately it is not easy to prove that the integral over the semi-circle vanishes in the limit $R \rightarrow \infty$, which is necessary to go back to the real-line integral. But since the final result will be shown to coincide with a result found by Grosche this point is not stressed here. The Green's function becomes

$$G(x'', x', E) = \frac{1}{\pi} \sinh(2\pi\sqrt{E}) |\Gamma(i\sqrt{E})|^2 e^{-(x'+x'')/2} \times W_{1/2, i\sqrt{E}}(e^{x'}) W_{1/2, i\sqrt{E}}(e^{x''}) \quad (69)$$

The connection of the Green's function to the energy eigenfunctions is given by Eq. (62). By the property $W_{\mu, \nu} = W_{\mu, -\nu}$ of the Whittaker function Eq. (69) is already real, so the real part need not be taken and the expression (69) is equal to

$$\pi \psi_E(x'') \psi_E^*(x') \rho(E)$$

From the general discussion earlier it is known that $\rho(E)$ must be identically 1 for δ -orthonormalized wave functions. This can be verified directly. Assume $\rho(E) = 1$ then the energy eigenfunctions are

$$\psi_E(x) = \left(\frac{\sinh(2\pi\sqrt{E})}{2\pi^2} \right)^{1/2} \Gamma(i\sqrt{E}) e^{-x/2} W_{1/2, i\sqrt{E}}(e^x) \quad (70)$$

and they are δ -orthonormalized. To see this use equations III.22 and III.23 of [46], which for the case at hand show that $\int_0^\infty u^{-2} W_{1/2, i\lambda}(u) W_{1/2, i\lambda'}(u) du = 2\pi \left| \frac{\Gamma(2i\lambda)}{\Gamma(i\lambda)} \right|^2 \delta(\lambda - \lambda')$. Further use $\Gamma(iy)\Gamma(-iy) = \pi y^{-1} [\sinh(\pi y)]^{-1}$ for real y ([47], 6.1.29) and thus

$$\begin{aligned} & \int_{-\infty}^{\infty} \psi_E(x) \psi_{E'}^*(x) dx \\ &= \frac{1}{2\pi^2} [\sinh(2\pi\sqrt{E}) \sinh(2\pi\sqrt{E'})]^{1/2} |\Gamma(i\sqrt{E})| |\Gamma(-i\sqrt{E'})| \\ & \quad \times \int_{-\infty}^{\infty} dx e^{-x} W_{1/2, i\lambda}(e^x) W_{1/2, i\lambda'}(e^x) \\ &= \frac{1}{2\pi^2} \sinh(2\pi\sqrt{E}) 2\pi |\Gamma(2i\sqrt{E})|^2 \delta(\sqrt{E} - \sqrt{E'}) = \delta(E - E') \end{aligned}$$




Figure 2: Morse Potential $\frac{1}{4}(\exp\{2x\} - 2\exp\{x\})$

Grosche [46] Fourier transforms Eq. (68) and thus gets the Feynman Kernel from which he reads off the eigenfunctions in momentum representation ($E = \lambda^2$ since $m = 1/2$ was chosen):

$$\psi_\lambda(x) = \left(\frac{\lambda \sinh(2\pi\lambda)}{\pi^2}\right)^{1/2} \Gamma(i\lambda) e^{-x/2} W_{1/2, i\lambda}(e^x) \quad (71)$$

He also proves that they are δ -orthonormalized wave-functions. Since $E = \lambda^2$, the momentum wave-functions ψ_λ are connected to the energy wave-functions ψ_E by $\int \psi_{\lambda'}^*(x) \psi_\lambda(x) dx = \delta(\lambda - \lambda') = \delta(\sqrt{E} - \sqrt{E'}) = 2\sqrt{E} \delta(E - E') = \int \psi_{E'}^*(x) \psi_E(x) \cdot 2\sqrt{E} dx$ (change effected by δ -function rule) and $\int \psi_\lambda(x'') \psi_\lambda^*(x') d\lambda = \delta(x'' - x') = \int \psi_E(x'') \psi_E^*(x') dE$ (change effected by the measure).

It was pointed out that, from a mathematical point of view, a density $\rho = 1$ followed from the δ -orthonormalization and the fact that the spectral family $[\mathbb{E}(\lambda)]$ is a family of projection operators. From a physical point of view, the density $\rho(E)$ can be interpreted as a measure for the degeneracy of the energy-wavefunctions. For the Morse potential $\frac{1}{4}(\exp\{2x\} - 2\exp\{x\})$ [see figure 2] one would expect no degeneracy or in other words $\rho(E) = 1$.

Since the Whittaker function $W_{1/2,0}(z) = e^{-z/2} z^{1/2}$ [as can easily be seen from the definition (67)], the x -dependence of $\psi_{E=0}(x)$ is $e^{-e^x/2}$. The rescaling factor is thus best determined with the choice $x'' = x' = b = -\infty$ where this function is 1. For small E the function $f_{-\infty, -\infty}(E) = \psi_E(-\infty) \psi_E^*(-\infty) \rho(E) \approx \pi^{-1} E^{-1/2}$ because $\sinh(2\pi\sqrt{E}) \approx 2\pi\sqrt{E}$, and $|\Gamma(i\sqrt{E})|^2 \approx 1/E$, and $\rho(E) = 1$. Inserting this E -dependence into the general formula (60) [set $a = -1/2$ there] one finds the inverse rescaling

factor

$$\int_0^\infty e^{-\nu TE} f_{-\infty, -\infty}(E) dE = (\pi \nu T)^{-1/2} \quad (72)$$

Because of the connection between the “Morse”-level and the original problem [Eqs. (64) and (65)] the rescaling factor, computed on the “Morse”-level, is valid for the original problem as well. Nonetheless it is useful to write down the problem and its solution on the “A”-level.

The eigenfunctions for A are²⁹:

$$\begin{aligned} A\psi_{E,\phi}(p, q) &:= A\langle U(p, q)Q^{1/2}\psi'_E|\phi\rangle = \langle U(p, q)Q^{1/2}H\psi'_E|\phi\rangle \\ &= \langle U(p, q)Q^{1/2}U^{-1}UHU^{-1}U\psi'_E|\phi\rangle \\ &= \langle U(p, q)Q^{1/2}U^{-1}H_{Morse}\psi_E|\phi\rangle \\ &= \langle U(p, q)Q^{1/2}U^{-1}E\psi_E|\phi\rangle = E\langle U(p, q)Q^{1/2}U^{-1}\psi_E|\phi\rangle \\ &= E\langle U(p, q)Q^{1/2}\psi'_E|\phi\rangle = E\psi_{E,\phi}(p, q) \end{aligned}$$

As was stated above the x -dependence of $\psi_{E=0}$ is $e^{-e^x/2}$. The inverse of the unitary transformation $\tilde{U}$ [Eq. (65)] is $\tilde{U}^{-1} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}^+)$, $\psi'(x) = (\tilde{U}^{-1}\psi)(x) = x^{-1/2}\psi(\ln x)$ and applying this to $\psi_{E=0}$ one has

$$\psi'_0(x) \propto x^{-1/2}e^{-\beta x}$$

Finally, by multiplying with $x^{1/2}$, one obtains:

$$e^{-\beta x}$$

This is exactly the minimum uncertainty state η_β for $\beta = 1/2$ and the result is no surprise, since this means that the eigenfunctions of A for eigenvalue 0 are exactly $\psi_{0,\phi}(p, q) = \langle pq|\phi\rangle$.

Observe that overall factors (normalizations) are completely unimportant since a quotient is formed anyway [compare e.g. Eq. (73)]. Even if they did “survive” it would not matter since a reproducing kernel multiplied with a factor is just as good a kernel (only the inner product in the reproducing kernel Hilbert space has to be redefined).

²⁹Observe that strictly $\langle U(p, q)Q^{1/2}\psi'_E|$ is a linear functional on the Hilbert space $L^2(\mathbb{R}^+)$, but the corresponding “ket” is not in the Hilbert space. But this “ket” is much better behaved than e.g. $|x\rangle$ which usually is used without the slightest hesitation. The $\psi_{E,\phi}(p, q)$ are not in $L^2(M_+)$, but are well-behaved functions (among them is e.g. the coherent state overlap), and they can be taken as generalized eigenvectors.

With an orthonormal basis ($|\phi_m\rangle$) in $L^2(\mathbb{R}^+)$ one can write the corresponding eigenfunctions of A as ψ_{E,ϕ_m} , and again by the above: $\psi_{0,\phi_m} = \langle pq|\phi_m\rangle$. Then use $\sum_m \int \psi_{E,\phi_m}(p,q)\psi_{E,\phi_m}^*(p',q')\rho(E)dE = \delta(p-p')\delta(q-q')$ [and still $\rho(E) = 1$] to find:

$$\begin{aligned}
& \lim_{\nu \rightarrow \infty} K_\nu e^{-\nu T A} \delta(p-p')\delta(q-q') \\
&= \lim_{\nu \rightarrow \infty} \frac{\sum_m \int_0^\infty e^{-\nu T E} \psi_{E,\phi_m}(p,q)\psi_{E,\phi_m}^*(p',q')\rho(E)dE}{\sum_{m'} \int_0^\infty e^{-\nu T E} \psi_{E,\phi_{m'}}(0,1)\psi_{E,\phi_{m'}}^*(0,1)\rho(E)dE} \\
&= \sum_m \psi_{0,\phi_m}(p,q)\psi_{0,\phi_m}^*(p',q') \\
&= \sum_m \langle pq|\phi_m\rangle \langle \phi_m|p'q'\rangle = \langle pq|p'q'\rangle
\end{aligned} \tag{73}$$

where $K_\nu^{-1} = e^{-\nu T A} \delta(p)\delta(q-1)|_{p=0,q=1} = (\pi\nu T)^{-1/2}$ by Eq. (72).

This is true at least in a distributional sense. Pointwise convergence on the other hand is not a problem, since the functional form of the overlap is known and is obviously continuous.

The last step is to formulate the weak coherent state path integral. Since the Feynman-Kac-Stratonovich representation of the operator $e^{-\nu T A}$ is already known from section (2.7), this is quickly done:

$$\begin{aligned}
& \lim_{\nu \rightarrow \infty} K_\nu e^{-\nu T A} \delta(p-p')\delta(q-q')|_{p=p'',q=q''} \\
&= \lim_{\nu \rightarrow \infty} K_\nu \mathcal{N}_\nu \int \exp\{-i \int q \dot{p} dt\} \exp\{-\frac{1}{2\nu} \int [\beta^{-1} q^2 \dot{p}^2 + \beta q^{-2} \dot{q}^2] dt\} \mathcal{D}p \mathcal{D}q \\
&= \lim_{\nu \rightarrow \infty} K_\nu \int \exp\{-i \int q dp\} d\mu_W^\nu \\
&= \langle p''q''|p'q'\rangle
\end{aligned} \tag{74}$$

The rescaling factor is $K_\nu = (\pi\nu T)^{1/2}$ and the Wiener measure $d\mu_W^\nu$ is the same as in Eq. (52). This is the sought-for weak coherent state path integral for $\beta = 1/2$ when the Hamiltonian vanishes.

Direct derivation

In the foregoing the weak coherent state path integral for $\beta = 1/2$ and a vanishing Hamiltonian was established. Its derivation relied on the general spectral approach. Since the (generalized) eigenfunctions of the

Morse operator are known, and because they are connected to the eigenfunctions of A , the isolating procedure can be developed step by step, i.e. $\lim_{\nu \rightarrow \infty} K_\nu e^{-\nu T A} \delta(p - p') \delta(q - q')$ can be computed directly. This will not lead to new knowledge, but since Eq. (74) is a central result it is good to confirm it:

$$\begin{aligned}
& \lim_{\nu \rightarrow \infty} K_\nu e^{-\nu T A} \delta(p - p') \delta(q - q') \\
&= \lim_{\nu \rightarrow \infty} \frac{\sum_m \int_0^\infty e^{-\nu T E} \psi_{E, \phi_m}(p, q) \psi_{E, \phi_m}^*(p', q') \rho(E) dE}{\sum_{m'} \int_0^\infty e^{-\nu T E} \psi_{E, \phi_{m'}}(0, 1) \psi_{E, \phi_{m'}}^*(0, 1) \rho(E) dE} \\
&= \lim_{\nu \rightarrow \infty} \frac{\sum_m \int_0^\infty e^{-\nu T E} \langle U(p, q) Q^{1/2} \psi'_E | \phi_m \rangle \langle \phi_m | U(p', q') Q^{1/2} \psi'_E \rangle \rho(E) dE}{\sum_{m'} \int_0^\infty e^{-\nu T E} \langle Q^{1/2} \psi'_E | \phi_{m'} \rangle \langle \phi_{m'} | Q^{1/2} \psi'_E \rangle \rho(E) dE} \\
&= \lim_{\nu \rightarrow \infty} \left\{ \int_0^\infty e^{-\nu T E} \int_0^\infty dx (qq')^{-1/2} e^{-ix(p-p')} [(x/q)^{1/2} \psi'_E(x/q)]^* \right. \\
&\quad \times [(x/q')^{1/2} \psi'_E(x/q')] \rho(E) dE \} \\
&\quad \times \left\{ \int_0^\infty e^{-\nu T E} \int_0^\infty dx [x^{1/2} \psi'_E(x)]^* [x^{1/2} \psi'_E(x)] \rho(E) dE \right\}^{-1} \quad (75)
\end{aligned}$$

Again $\rho(E) = 1$ since the eigenfunctions are δ -orthonormalized. The denominator of Eq. (75) will be examined first and it will be more convenient to work with the eigenfunctions ψ_λ . The connection between ψ_λ and ψ_E was demonstrated in Eqs. (71), (70). Recall that, by definition, $\psi'_\lambda(x) = U^{-1} \psi_\lambda(x)$ with $U^{-1} : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R}^+)$, $\psi'(x) = (U^{-1} \psi)(x) = x^{-1/2} \psi(\ln x)$. With $\beta = 1/2$ one has:

$$x^{1/2} \psi'_\lambda(x) = x^{1/2} x^{-1/2} \psi_\lambda(\ln(x)) = \left(\frac{\lambda \sinh(2\pi\lambda)}{\pi^2} \right)^{1/2} \Gamma(i\lambda) x^{-1/2} W_{1/2, i\lambda}(x)$$

For the x -integration, III.22 of Grosche [46] [or [45] page 858] is used:

$$\begin{aligned}
& \int_0^\infty x^{\rho-1} W_{\kappa, \mu}(x) W_{\lambda, \nu}(x) dx \\
&= \frac{\Gamma(1+\mu+\nu+\rho) \Gamma(1-\mu+\nu+\rho) \Gamma(-2\nu)}{\Gamma(1/2-\lambda-\nu) \Gamma(3/2-\kappa+\nu+\rho)} \\
&\quad \times {}_3F_2(1+\mu+\nu+\rho, 1-\mu+\nu+\rho, 1/2-\lambda+\nu; \\
&\quad \quad 1+2\nu, 3/2-\kappa+\nu+\rho; 1) \\
&\quad + \frac{\Gamma(1+\mu-\nu+\rho) \Gamma(1-\mu-\nu+\rho) \Gamma(2\nu)}{\Gamma(1/2-\lambda+\nu) \Gamma(3/2-\kappa-\nu+\rho)} \\
&\quad \times {}_3F_2(1+\mu-\nu+\rho, 1-\mu-\nu+\rho, 1/2-\lambda-\nu; \\
&\quad \quad 1-2\nu, 3/2-\kappa-\nu+\rho; 1)
\end{aligned}$$

Using $\rho = 0$, $\kappa = \lambda = 1/2$, $\mu = -\nu = i\lambda$, the equation becomes:

$$\begin{aligned}
& \int_0^\infty x^{-1} W_{1/2, i\lambda}(x) W_{1/2, i\lambda}(x) dx \\
&= \frac{\Gamma(1)\Gamma(1-2i\lambda)\Gamma(2i\lambda)}{\Gamma(i\lambda)\Gamma(1-i\lambda)} \\
&\quad \times {}_3F_2(1, 1-2i\lambda, -i\lambda; 1-2i\lambda, 1-i\lambda; 1) \\
&\quad + \frac{\Gamma(1+2i\lambda)\Gamma(1)\Gamma(-2i\lambda)}{\Gamma(-i\lambda)\Gamma(1+i\lambda)} \\
&\quad \times {}_3F_2(1+2i\lambda, 1, i\lambda; 1+2i\lambda, 1+i\lambda; 1)
\end{aligned}$$

With $\Gamma(1+x) = x\Gamma(x)$ this relation simplifies:

$$\begin{aligned}
& 2 \frac{\Gamma(2i\lambda)\Gamma(-2i\lambda)}{\Gamma(i\lambda)\Gamma(-i\lambda)} \\
& [{}_3F_2(1, 1-2i\lambda, -i\lambda; 1-2i\lambda, 1-i\lambda; 1) \\
& + {}_3F_2(1+2i\lambda, 1, i\lambda; 1+2i\lambda, 1+i\lambda; 1)]
\end{aligned}$$

Observe that by $\Gamma(iy)\Gamma(-iy) = \frac{\pi}{y \sinh(\pi y)}$ [e.g. [47], 6.1.29, p. 256], the prefactor cancels the normalization of the wave functions, except for a factor π .

Now, using the definition of the generalized hypergeometric function (or Barnes's extended hypergeometric function)

$${}_pF_q(\alpha_1, \dots, \alpha_p; \beta_1, \dots, \beta_q; z) := \sum_{n=0}^{\infty} \frac{(\alpha_1)_n \dots (\alpha_p)_n}{(\beta_1)_n \dots (\beta_q)_n} \frac{z^n}{n!}$$

with $(\alpha)_n = \Gamma(\alpha+n)/\Gamma(\alpha)$ (Gegenbauer coefficients), one can proceed formally

$$\begin{aligned}
& [{}_3F_2(1, 1-2i\lambda, -i\lambda; 1-2i\lambda, 1-i\lambda; 1) \\
& + {}_3F_2(1+2i\lambda, 1, i\lambda; 1+2i\lambda, 1+i\lambda; 1)] \\
&= [{}_2F_1(1, -i\lambda; 1-i\lambda; 1) \\
& + {}_2F_1(1, i\lambda; 1+i\lambda; 1)] \\
&= \sum_{n=0}^{\infty} \frac{-i\lambda}{-i\lambda+n} + \sum_{n=0}^{\infty} \frac{i\lambda}{i\lambda+n} \\
&=: \sum_{n=0}^{\infty} \frac{2\lambda^2}{n^2 + \lambda^2} = \pi\lambda \coth(\pi\lambda) + 1
\end{aligned}$$

It is important to notice that the first and second equality are merely symbolic since the expressions are undefined (complex infinite terms). Because $|z| = 1$ is the radius of convergence for the hypergeometric series defining ${}_2F_1(a, b; c; z)$ ([47], 15.1.1), only conditional convergence exists for $-1 < \Re(c - a - b) \leq 0$ (the point $z = 1$ is even excluded). Meaning was given to the undefined sums of infinite terms by setting up an ordering prescription, thus arriving at the last line. The same ordering prescription arises if one replaces the argument 1 by $\delta < 1$, sums the now absolutely convergent series and finally takes the limit $\delta \rightarrow 1$.

The complete denominator has become

$$\pi^{-1} \int_0^\infty e^{-\nu T \lambda^2} [\pi \lambda \coth(\pi \lambda) + 1] d\lambda =: h(\nu T)$$

This integral exists and $h(\nu T)$ is well-defined. Using the power-series expansion for the hyperbolic tangent the result is:

$$h(\nu T) = \pi^{-1/2} (\nu T)^{-1/2} [1 + \pi^2 / (12 \nu T) + O((\nu T)^{-2})] \quad (76)$$

The first term in the expansion is equal to the inverse rescaling factor computed in Eq. (72).

There is yet another way to treat the denominator. The idea is to extract the λ -dependence by series expansions, but the calculations are more complicated and so this part is presented in appendix (A.4).

The remaining task is to compute (again the λ -notation is chosen rather than the E -notation):

$$\begin{aligned} & \lim_{\nu \rightarrow \infty} \pi^{1/2} \sqrt{\nu T} \left(1 - \frac{\pi^2}{12\nu} + O(\nu^{-2}) \right) \\ & \times \int_0^\infty d\lambda e^{-\nu T \lambda^2} \int_0^\infty dx (qq')^{-1/2} e^{-ix(p-p')} [(x/q)^{1/2} \psi'_\lambda(x/q)]^* \\ & \times [(x/q')^{1/2} \psi'_\lambda(x/q')] \end{aligned}$$

First, notice that for $p = p'$ and $q = q'$, the x -integral will be the same as the one encountered in the denominator (make the transformation $x \rightarrow xq$) and thus the whole expression is exactly 1 [which can already be seen by symbolic manipulation in the third line of Eq. (75).]. This result was expected, since $\langle pq|pq \rangle = 1$.

Unfortunately it is very hard (if not impossible) to find an exact solution for the x -integral.

Even for $q = q'$ and after expanding $e^{-ix(p-p')}$ into a power series, it does not seem possible to use the integration formula for the Whittaker functions and powers of x and to sum things up using the expansions of the generalized hypergeometric functions, since this does not converge. However, one still expects the integrals to exist. This is a pity, since varying p and p' alone would be enough, since the functions $\langle pq|p'q \rangle$ already span the space.

But one can simply look at $[(x/q)^{1/2}\psi'_\lambda(x/q)]^*[(x/q')^{1/2}\psi'_\lambda(x/q')]$ and write it as $f_{x,q,q'}(\lambda)$. This is continuous in λ , and one can write $f_{x,q,q'}(\lambda) = g_{x,q,q'} + O_{x,q,q'}(\lambda)$. The λ -independent function g is exactly $[(x/q)^{1/2}\psi'_0(x/q)]^*[(x/q')^{1/2}\psi'_0(x/q')] = 2\pi^{-1}\eta(x/q)^*\eta(x/q')$, and the λ - and x -integrals will factorize. The first integral yields $\frac{1}{2}\pi^{1/2}(\nu T)^{-1/2}$, while the second integral gives $\pi^{-1}\langle pq|p'q' \rangle$. So, the limit of diverging diffusion constant yields the desired reproducing kernel.

One has to make sure that, for the higher orders $O_{x,q,q'}(\lambda)$, the x -integrals exist. Otherwise the expansion followed by interchange of summation and integration will not be justified. But this proof is simple since by the Cauchy-Schwarz inequality

$$\begin{aligned} & \left| \int_0^\infty dx [q^{-1/2}e^{ixp}(x/q)^{1/2}\psi'_{\lambda^2}(x/q)]^* [q'^{-1/2}e^{ixp'}(x/q')^{1/2}\psi'_{\lambda^2}(x/q')] \right|^2 \\ & \leq \int_0^\infty dx [q^{-1/2}e^{ixp}(x/q)^{1/2}\psi'_{\lambda^2}(x/q)]^2 \\ & \quad \times \int_0^\infty dx [q'^{-1/2}e^{ixp'}(x/q')^{1/2}\psi'_{\lambda^2}(x/q')]^2 \\ & = q^{1/2} \int_0^\infty dx [x^{1/2}\psi'_{\lambda^2}(x)]^2 \cdot q'^{1/2} \int_0^\infty dx [x^{1/2}\psi'_{\lambda^2}(x)]^2 \end{aligned}$$

and the existence of the right-hand side has already been proved. These integrals are like the ones in the denominator (compare above). Once this is established, the exact result of any of these integrations (which might be very hard to determine) is of no consequence since integration over λ gives in general ($n \geq 1$):

$$\int_0^\infty e^{-\nu T \lambda^2} \lambda^n d\lambda = 1/\sqrt{\nu T} \cdot \Gamma((n+1)/2)(\nu T)^{-n/2}$$

Whatever the corresponding x -integral is, these terms will all vanish in the limit $\nu \rightarrow \infty$.

In this way, the isolating procedure has been directly computed and the results (73) and (74) are confirmed.

3.2.2 Introducing dynamics

Since the only case in which the spectral approach worked was $\beta = 1/2$, this value is assumed throughout this subsection. Dynamics are introduced by the quantum Hamiltonian $\mathcal{H}$, which is a function of the basic kinematical operators Q and D . The goal is to represent the propagator $\langle p'' q'' | \exp\{-iT\mathcal{H}\} | p' q' \rangle$ as a weak coherent state path integral.

Klauder [27] proposed

$$\begin{aligned}
& \langle p'' q'' | \exp\{-iT\mathcal{H}\} | p' q' \rangle \\
&= \lim_{\nu \rightarrow \infty} K_\nu \mathcal{N}_\nu \int \exp\{-i \int [q\dot{p} + h_w(p, q)] dt\} \\
&\quad \times \exp\{-\frac{1}{2\nu} \int [\beta^{-1} q^2 \dot{p}^2 + \beta q^{-2} \dot{q}^2] dt\} \mathcal{D}p \mathcal{D}q \\
&= \lim_{\nu \rightarrow \infty} K_\nu \int \exp\{-i \int [q dp + h_w(p, q) dt]\} d\mu_W^\nu \tag{77}
\end{aligned}$$

as the path integral for a class of Hamiltonians which contains at least all Hamiltonians polynomial in Q and D . The new symbol $h_w(p, q)$, interpreted as the classical Hamiltonian associated with the quantum Hamiltonian, is implicitly given by

$$\langle p'' q'' | \mathcal{H} | p' q' \rangle = \lim_{\nu \rightarrow \infty} K_\nu \int \exp\{-i \int q dp\} T^{-1} \int h_w(p, q) dt d\mu_W^\nu \tag{78}$$

and will be called the weak symbol.

The whole conjecture is based on the observation that, for a linear Hamiltonian $RQ + SD$, the propagator can be reduced to a mere overlap. This can be seen as follows: Write $\exp\{-it[RQ + SD]\} = \exp\{-iF(t)Q\} \exp\{-iG(t)D\}$. Differentiation leads to the differential equation $RQ + SD = (\dot{F} + F\dot{G})Q + \dot{G}D$, where $\exp\{-iFQ\}D \exp\{iFQ\} = D + FQ$ was used at an intermediate step. The solution of this equation with the initial values $F(0) = G(0) = 0$ is $F(t) = R/S(1 - \exp\{-St\})$ and $G(t) = St$. With this, one has

$$\begin{aligned}
& e^{i \ln q'' D} e^{-i p'' Q} e^{-i(RQ+SD)T} = e^{i \ln q'' D} e^{-i p'' Q} e^{-i R/S(1-e^{ST})Q} e^{-i STD} \\
& = e^{i \ln q'' D} e^{-i STD} e^{i STD} e^{-i[p''+R/S(1-e^{ST})]Q} e^{-i STD} \\
& = e^{i \ln(q'' e^{-ST})D} e^{-i e^{ST}[p''+R/S(1-e^{ST})]Q} \\
& = e^{i \ln(q'' e^{-ST})D} e^{-i[p'' e^{ST}+R/S(e^{ST}-1)]Q}
\end{aligned}$$

and thus

$$\langle p'' q'' | e^{-i(RQ+SD)T} | p' q' \rangle = \langle p'' e^{ST} + R/S \cdot (e^{ST} - 1), q'' e^{-ST} | p' q' \rangle \quad (79)$$

Consequently, the problem is already solved for a linear Hamiltonian and it remains to determine the weak symbol associated with $\mathcal{H} = RQ + SD$. According to Eq. (74) the path integral for this is

$$\lim_{\nu \rightarrow \infty} K_\nu \int_{p', q'}^{p'' e^{ST} + R/S \cdot (e^{ST} - 1), q'' e^{-ST}} \exp\{-i \int q dp\} d\mu_W^\nu \quad (80)$$

Since this is a well-defined functional integral, one can change variables

$$\begin{aligned}
p(t) & \rightarrow p(t) e^{St} + R/S(e^{St} - 1) \\
q(t) & \rightarrow q(t) e^{-St}
\end{aligned}$$

and the integrand becomes now $\exp\{-i \int (q e^{-St} d[p e^{St} + R/S(e^{St} - 1)])\} = \exp\{-i \int [q dp + (Rq + Spq)dt]\}$. The new measure is³⁰

$$\begin{aligned}
d\tilde{\mu}_W^\nu & = \mathcal{N}_\nu \exp\{-\frac{1}{2\nu} \int [\beta^{-1} (q e^{-St})^2 (p e^{St} + \frac{R}{S}(e^{St} - 1))^{\bullet 2} \\
& \quad + \beta (q e^{-St})^{-2} (q e^{-St})^{\bullet 2}] dt\} \mathcal{D}[p e^{St} + \frac{R}{S}(e^{St} - 1)] \mathcal{D}(q e^{-St})
\end{aligned}$$

But, since the symbolic measure is actually

$$\begin{aligned}
\mathcal{D}[p e^{St} + \frac{R}{S}(e^{St} - 1)] & = \lim_{\epsilon \rightarrow 0} \prod_{k=1}^N d[p(t) e^{St}|_{t=k\epsilon} + \frac{R}{S}(e^{St} - 1)|_{t=k\epsilon}] \\
& = \lim_{\epsilon \rightarrow 0} \prod_{k=1}^N [dp(t) e^{St}|_{t=k\epsilon} + (p S e^{St} + R e^{St}) dt|_{t=k\epsilon}] \\
& = \lim_{\epsilon \rightarrow 0} \prod_{k=1}^N [dp_k e^{Sk\epsilon} + (p_k S e^{Sk\epsilon} + R e^{Sk\epsilon}) \epsilon] \\
& = \lim_{\epsilon \rightarrow 0} \prod_{k=1}^N dp_k e^{Sk\epsilon} = \mathcal{D}p \prod_t e^{St}
\end{aligned}$$

³⁰(...)[•] means the time derivative of the expression in parentheses

and analogous $\mathcal{D}(qe^{-St}) = \mathcal{D}q \prod_t e^{-St}$, the new measure can be expressed in terms of the old one as

$$\begin{aligned} d\tilde{\mu}_W^\nu &= \exp\left\{-\frac{1}{2\nu}\int[\beta^{-1}q^2((Sp+R)^2+2(Sp+R)\dot{p})\right. \\ &\quad \left.+\beta q^{-2}(S^2q^2-2Sq\dot{q})]dt\right\} d\mu_W^\nu \\ &= \exp\left\{-\frac{1}{2\nu}\int[\beta^{-1}q^2((Sp+R)^2dt+2(Sp+R)dp)\right. \\ &\quad \left.+\beta q^{-2}(S^2q^2dt-2Sqdq)]\right\} d\mu_W^\nu \end{aligned}$$

The first equality is again formal and gains meaning by the last line, where the stochastic integrals have to be understood in the Stratonovich sense as usual. The change of variables has introduced additional terms in the exponent of the formal expression which are at most linear in $\dot{p}$ or $\dot{q}$ respectively. These terms are not critical, since, in the limit of diverging diffusion constant ν , they will vanish. This means that the total change of the measure disappears in the limit. Thus, one can write the path integral with the old measure $d\mu_W^\nu$ (instead of with the new $d\tilde{\mu}_W^\nu$), and one finds

$$\langle p''q''|e^{-i(RQ+SD)T}|p'q'\rangle = \lim_{\nu\rightarrow\infty} K_\nu \int_{p',q'}^{p'',q''} \exp\{-i\int[qdp+(Rq+Spq)dt]\} d\mu_W^\nu \quad (81)$$

Now, the weak symbol can be read off:

$$h_w(p, q) = Rq + Spq \quad (82)$$

The generalization to other Hamiltonians is based on the linearity, completeness and irreducibility of the basic operators Q and D by virtue of which $\lim_{J\rightarrow\infty} \sum_{j=1}^J \alpha_j e^{-i(R_j Q + S_j D)}$ weakly converges to any (bounded) operator such as $e^{-i\mathcal{H}T}$. Thus,

$$\begin{aligned} \langle p''q''|e^{-i\mathcal{H}T}|p'q'\rangle &= \lim_{J\rightarrow\infty} \langle p''q''|\sum_{j=1}^J \alpha_j e^{-i(R_j Q + S_j D)}|p'q'\rangle \\ &= \lim_{J\rightarrow\infty} \lim_{\nu\rightarrow\infty} K_\nu \int \exp\{-i\int q dp\} \\ &\quad \times [\sum_{j=1}^J \alpha_j \exp\{-i\int (R_j q + S_j pq)dt\}] d\mu_W^\nu \quad (83) \end{aligned}$$

and the question, on which the next steps depend, is: can the two limits be interchanged? In spite of some effort this question is not yet answered. Assuming that they can, however, one obtains

$$\begin{aligned}
& \langle p'' q'' | e^{-i\mathcal{H}T} | p' q' \rangle \\
&= \lim_{\nu \rightarrow \infty} K_\nu \int e^{-i \int q dp} \left[\lim_{J \rightarrow \infty} \sum_{j=1}^J \alpha_j e^{-i \int (R_j q + S_j p q) dt} \right] d\mu_W^\nu \quad (84)
\end{aligned}$$

The expression $[\lim_{J \rightarrow \infty} \sum_{j=1}^J \alpha_j e^{-i \int (R_j q + S_j p q) dt}] =: F[\int q dt, \int p q dt]$ is, unfortunately, not of the form $e^{-i \int h_w(p, q) dt}$ for a general, local Hamiltonian h_w , e.g. $e^{-i \int q^2 dt}$ with Hamiltonian q^2 . To produce local Hamiltonians, one would need distributions $R(t)$ and $S(t)$ instead of the constants R and S . Then, taking e.g. $R(t) = \delta(t - \tau)$, one gets a local expression $q(\tau)$ and, by forming functions thereof, local Hamiltonians. This was proposed in [27]. However, the construction of distributions from piecewise constant functions would require yet another limiting process, and, again, the interchangeability of the limits is questionable.

In the case of a linear Hamiltonian, the weak symbol was shown to be $h_w(p, q) = Rq + Spq$. This is exactly what one would expect since the connection of the basic operators Q and D to classical variables is, according to the weak correspondence principle, q and pq , respectively. But, the correspondence for a more general Hamiltonian is not immediately clear and remains to be determined.

3.2.3 Regularizing approach

In the second to last subsection it could be seen that a rescaling procedure to isolate the reproducing kernel Hilbert space can - in the spectral approach - only work in the case $\beta = 1/2$. To proceed to the final goal - a well-defined weak coherent state path integral - in the case $0 < \beta < 1/2$ ³¹ one has to discuss new methods.

DKP [26] were able to use the full power of functional analysis and its operator techniques to prove the result in the case $\beta > 1/2$ since the problem was confined to the Hilbert space $L^2(M_+)$. Instead of trying to work outside of Hilbert space, as was done in the previous two subsections, one can try to bring the problem back into Hilbert space by introducing a new regularization parameter ε which will go to zero in the very end. Two distinct proposals will be considered.

The first idea is to reestablish the fiducial vector admissibility condition $\langle Q^{-1} \rangle = \int_0^\infty x^{-1} |\eta_\beta(x)|^2 dx < \infty$, which was important to ensure a resolution

³¹or in other words, since the (scalar) curvature of the Lobachevsky half-plane is $R = -2/\beta$: for a higher negative curvature than -4

of unity and which failed in the case $\beta \leq 1/2$ since then $\int_0^\infty x^{-1} x^{2\beta-1} e^{-2\beta x} = \infty$ due to lack of integrability at 0. The regularization factor $e^{-\varepsilon/x}$ takes care of that leading to a modified fiducial vector $\eta_{\beta,\varepsilon} = N_{\beta,\varepsilon} x^{\beta-1/2} e^{-\beta x} e^{-\varepsilon/x}$ ³². The affine group acts on this new fiducial vector producing the group-defined coherent states $|pq\rangle_\varepsilon$. Their overlap is now

$$\begin{aligned} & \langle pq|rs\rangle_\varepsilon \\ &= \int_0^\infty dx (e^{ipx} e^{-\ln(q)(x\partial_x+1/2)} N_{\beta,\varepsilon} x^{\beta-1/2} e^{-\beta x} e^{-\varepsilon/x})^* \\ & \quad \times (e^{irx} e^{-\ln(s)(x\partial_x+1/2)} N_{\beta,\varepsilon} x^{\beta-1/2} e^{-\beta x} e^{-\varepsilon/x}) \\ &= |N_{\beta,\varepsilon}|^2 (qs)^{-1/2} \\ & \quad \times \int_0^\infty dx e^{-i(p-r)x} \left(\frac{x}{q}\right)^{\beta-1/2} \left(\frac{x}{s}\right)^{\beta-1/2} e^{-\beta(q^{-1}+s^{-1})x} e^{-\varepsilon(q+s)/x} \end{aligned}$$

with $|N_{\beta,\varepsilon}|^2 = (\int_0^\infty dx x^{2\beta-1} e^{-2\beta x} e^{-2\varepsilon/x})^{-1}$

From the second to the third line $e^{-\alpha\partial_x}\phi(x) = \phi(x-\alpha)$ was used and thus $e^{-\ln(q)(x\partial_x)}\phi(x) = e^{-\ln(q)\partial_u}\phi(e^u) = \phi(e^{u-\ln(q)}) = \phi(e^{\ln(x)-\ln(q)}) = \phi(x/q)$.

A new operator B_ε which annihilates this overlap could be computed which would lead to a new operator A_ε .

But, there are reasons not to pursue this course. Retaining the group definition of the coherent states on the positive side, one is confronted with the problem that the group will no longer operate on an extremal weight vector (and minimum uncertainty state). Thus one can not expect any linear analog of the former $(Q - 1 + i\beta^{-1}D)|\eta_\beta\rangle = 0$ to hold³³. This has serious implications for the new path integral, i.e., the path integral with Wiener measure. It was the linear complex polarization that led to a first order differential operator B and thus to a second order differential operator $A = \frac{1}{2}\beta B^\dagger B$. It was due to this second order property that the Wiener measure emerged and the path integral became a well-defined functional integral [see 2.7]. Whereas it would still be possible to derive the old, standard path integral, the Wiener measure path integral can not exist for fiducial vectors which are not extremal weight vectors.

The second possibility is to regularize the overlap directly by a multiplicative ε -dependent factor which ensures square integrability on the Lobachevsky manifold. From the second to last subsection and appendix (A.3)

³²Observe that in the case $\beta = 1/2$ a weaker regularization, namely x^ε , would be enough, which could be interpreted as a deformation of the operator Q , thus leading to a deformed version of the affine group.

³³And one has to give up analyticity, which previously held apart from a factor $(qs)^{-\beta}$.

(where $0 \notin \text{spec}(A)$ for $\beta < 1/2$) was established) it is known that for $0 < \beta < 1/4$ one is forced to regularize both in p and q . So the case $1/4 < \beta < 1/2$, where no p -regularization is needed, will be examined first.

Case $1/4 < \beta \leq 1/2$

Let

$$\langle pq|rs \rangle_\varepsilon = N_\varepsilon \langle pq|rs \rangle e^{-(q+s)\varepsilon} \quad (85)$$

where N_ε is a normalization constant, which is w.l.o.g. chosen to be real. The extra factor $e^{-(q+s)\varepsilon}$ goes to 1 in the limit $\varepsilon \rightarrow 0$.

The modified overlap lies in the Hilbert space $L^2(M_+)$ and its norm is computed to be:

$$\begin{aligned} 1 &\stackrel{!}{=} N_\varepsilon^* N_\varepsilon \int_0^\infty dq (qs)^{-2\beta} e^{-2q\varepsilon} \int_{-\infty}^\infty dp [(q^{-1} + s^{-1})^2 + \beta^{-2}(p-r)^2]^{-2\beta} \\ &= N_\varepsilon^* N_\varepsilon \int_0^\infty dq (qs)^{-2\beta} e^{-2q\varepsilon} (q^{-1} + s^{-1})^{1-4\beta} \int_{-\infty}^\infty dp (1 + \beta^{-2}p^2)^{-2\beta} \\ &= N_\varepsilon^* N_\varepsilon \int_0^\infty dq \varepsilon^{-1} (qs)^{-2\beta} \varepsilon^{2\beta} e^{-2q} (\varepsilon q^{-1} + s^{-1})^{1-4\beta} \cdot I_p \\ &\leq N_\varepsilon^* N_\varepsilon I_p s^{2\beta-1} \varepsilon^{2\beta-1} \int_0^\infty dq q^{-2\beta} e^{-2q} \\ &= N_\varepsilon^* N_\varepsilon I_p s^{2\beta-1} \varepsilon^{2\beta-1} 2^{2\beta} \Gamma(1-2\beta) \end{aligned}$$

So $N_\varepsilon^* N_\varepsilon \geq \text{const} \cdot \varepsilon^{1-2\beta}$ and this is the correct asymptotic behavior. Instead of the explicit expression for the normalization constant one can use $\langle xy|xy \rangle_\varepsilon = N_\varepsilon e^{-2y\varepsilon}$ (for arbitrary $x \in \mathbb{R}$, $y \in \mathbb{R}^+$) and write $\langle pq|rs \rangle = \lim_{\varepsilon \rightarrow 0} \langle pq|rs \rangle_\varepsilon / \langle xy|xy \rangle_\varepsilon$ in a self-consistent way. For a shorter notation set

$$\langle xy|xy \rangle_\varepsilon =: c_{\beta,\varepsilon} \quad (86)$$

One could also use a different regularization like $e^{-(q^2+s^2)\varepsilon}$ or something similar. But, the expression $e^{-(q+s)\varepsilon}$ is simpler and thus the preferred choice.

Certainly, the new states $|pq\rangle_\varepsilon$ are not affine group-defined coherent states. The group law for the affine group in p - q notation is $(p, q) * (p', q') = (q^{-1}p' + p, qq')$. For the new $U_\varepsilon(p, q) = U(p, q)e^{-q\varepsilon}$, one would have to add q and q' in the extra factor instead of multiply them. Indeed, they are not

group defined at all³⁴, but it will shortly be shown that they still share certain properties of coherent states:

1) Continuity in the labels is no problem since joint continuity in the overlap is obvious.

2) The old $|pq\rangle$ spanned the space, i.e., any $|\psi\rangle$ could be expressed as $\sum_{n=0}^N \alpha_n |p_n q_n\rangle$ or Cauchy limits thereof. The new $|pq\rangle_\varepsilon$ still span the space since they are related to the old ones by $\sum \alpha_n |p_n q_n\rangle_\varepsilon = \sum \alpha_n N_\varepsilon^{1/2} e^{-q_n \varepsilon} |p_n q_n\rangle =: \sum \beta_n |p_n q_n\rangle$.

Therefore, these new $|pq\rangle_\varepsilon$ certainly are Klauder states. However, there is ambiguity as to whether they are coherent states or, more likely, non-group-defined weak coherent states. Since they are only a mathematical aid used in intermediate steps this point will not be clarified here.

The new operator B_ε which annihilates the modified kernel is derived by exploiting analyticity (up to a factor): $[(q^{-1} + s^{-1}) + i\beta^{-1}(p - r)]^{-2\beta} =: Y$ is analytic, so $\partial_{(q^{-1} - i\beta^{-1}p)} Y = (-\frac{1}{2}q^2 \partial_q + \frac{1}{2}i\beta \partial_p) Y = 0$. Writing Y as $e^{q\varepsilon}(qs)^\beta \langle pq|rs\rangle_\varepsilon$, one moves $e^{q\varepsilon}(qs)^\beta$ to the left of this operator. Then $e^{q\varepsilon}(qs)^\beta$ can be cancelled since this expression is everywhere nonzero. The result is the new operator

$$B_\varepsilon = (\beta^{-1}q\partial_q + \beta^{-1}q\varepsilon + 1 - iq^{-1}\partial_p)$$

for which $B_\varepsilon \langle pq|rs\rangle_\varepsilon = 0$.

Define $A_\varepsilon := \frac{1}{2}\beta B_\varepsilon^\dagger B_\varepsilon$, then

$$\begin{aligned} A_\varepsilon &= \frac{1}{2}\beta(-iq^{-1}\partial_p + 1 - \beta^{-1}\partial_q q + \beta^{-1}q\varepsilon)(-iq^{-1}\partial_p + 1 + \beta^{-1}q\partial_q + \beta^{-1}q\varepsilon) \\ &= \frac{1}{2}\{\beta[-iq^{-1}\partial_p + 1 + \beta^{-1}q\varepsilon]^2 - \beta^{-1}\partial_q q^2 \partial_q - 1 - 2\beta^{-1}q\varepsilon\} \end{aligned} \quad (87)$$

Here $1/4 < \beta \leq 1/2$ and p, q are the variables of the Lobachevsky half-plane M_+ ($q > 0$, metric: $g^{pp} = \beta q^{-2}$, $g^{qq} = \beta^{-1}q^2$, $g^{pq} = g^{qp} = 0$, Laplace-Beltrami operator: $\beta^{-1}\partial_q q^2 \partial_q + \beta q^{-2} \partial_p^2$).

Setting $-i\partial_p = P_p$, $p = Q_p$ (does not appear), $-i\partial_q = P_q$, and $q = Q_q$ and writing the metric with a capital G to reflect its operator character ($a, b \in \{p, q\}$), one finds:

³⁴For $\beta = 1/2$ a weaker regularization is enough, namely $q^{-\varepsilon} = e^{-\ln(q)\varepsilon}$, leading to group-defined coherent states with a modified dilation operator $D - i\varepsilon$ and thus to a modified affine group.

$$\begin{aligned}
A_\varepsilon &= \frac{1}{2}\{(P_a - \mathcal{A}_a)G^{ab}(P_b - \mathcal{A}_b) + V(Q_q, Q_p)\} \\
&= \frac{1}{2}\{\beta Q_q^{-2}(P_p - \mathcal{A}_p)^2 + \beta^{-1}(P_q - \mathcal{A}_q)Q_q^2(P_q - \mathcal{A}_q) + V(Q_q)\}
\end{aligned}$$

where

$$\begin{aligned}
\mathcal{A}_p &= -Q_q(1 + \beta^{-1}Q_q\varepsilon) \\
\mathcal{A}_q &= 0 \\
V(Q_q) &= -2\beta^{-1}Q_q\varepsilon - 1
\end{aligned}$$

Effectively the magnetic field has been changed (augmented by an ε -dependent part), which alters the vector potential $\mathcal{A} = (\mathcal{A}_p, \mathcal{A}_q)$. Additionally, the explicit potential V underwent a modification as well.

Next, define

$$K_{\nu, \varepsilon}^0(p'', p', q'', q') := e^{\nu T/2} \int \exp\{-i \int (q + \beta^{-1}q^2\varepsilon)dp + \frac{\nu}{2} \int \beta^{-1}qdt\varepsilon\} d\mu_W^\nu \quad (88)$$

with Wiener measure, pinned at (p', q') for $t = 0$ and (p'', q'') for $t = T$, [see Eq. (52)]:

$$\mathcal{N} \exp\{-\frac{1}{2\nu} \int [\beta^{-1}q^2\dot{p}^2 + \beta q^{-2}\dot{q}^2]dt\} \mathcal{D}p \mathcal{D}q = d\mu_W^\nu \quad (89)$$

This is the Feynman-Kac-Stratonovich representation of the kernel of the operator $\exp\{-\nu T A_\varepsilon\}$ and it is derived in the following way:

$$\begin{aligned}
& \exp\{-\nu T A_\varepsilon\} \delta(p-p') \delta(q-q')|_{p=p'', q=q''} \\
= & \exp\{-\tfrac{1}{2}\nu T [\beta(-iq^{-1}\partial_p + 1 + \beta^{-1}q\varepsilon)^2 - \beta^{-1}\partial_q q^2 \partial_q - 1 - 2\beta^{-1}q\varepsilon]\} \\
& \times \int e^{ix(p-p') - ik(q-q')} \frac{dx dk}{(2\pi)^2} |_{p=p'', q=q''} \\
= & e^{\nu T/2} \lim_{N \rightarrow \infty} [\exp\{-\tfrac{1}{2}\nu \delta [\beta(-iq^{-1}\partial_p + 1 + \beta^{-1}q\varepsilon)^2 - 2\beta^{-1}q\varepsilon]\} \\
& \times \exp\{-\tfrac{1}{2}\nu \delta (-\beta^{-1}\partial_q q^2 \partial_q)\}]^N \int e^{ix(p-p') - ik(q-q')} \frac{dx dk}{(2\pi)^2} |_{p=p'', q=q''} \\
= & \lim_{N \rightarrow \infty} e^{\nu T/2} \int \exp\{i \sum x_{l+1/2}(p_{l+1} - p_l) - ik_{l+1/2}(q_{l+1} - q_l)\} \\
& \times \exp\{-\tfrac{1}{2}\nu \delta \sum [\beta(q_l^{-1}x_{l+1/2} + 1 + \beta^{-1}q_l\varepsilon)^2 - 2\beta^{-1}q_l\varepsilon]\} \\
& \times \exp\{-\tfrac{1}{2}\nu \delta \sum \beta^{-1}k_{l+1/2}^2 q_l^2\} \prod_{l=0}^N \frac{dk_{l+1/2} dx_{l+1/2}}{(2\pi)^2} \prod_{l=1}^N dp_l dq_l \\
=: & e^{\nu T/2} \mathcal{N} \int \exp\{i \int (x\dot{p} - k\dot{q}) dt\} \\
& \times \exp\{-\tfrac{1}{2}\nu \int \{\beta(q^{-1}x + 1 + \beta^{-1}q\varepsilon)^2 - 2\beta^{-1}q\varepsilon + \beta^{-1}k^2 q^2\} dt\} \\
& \times \mathcal{D}x \mathcal{D}k \mathcal{D}p \mathcal{D}q \\
= & e^{\nu T/2} \mathcal{N} \int \exp\{i \int [(x - q - \beta^{-1}q^2\varepsilon)\dot{p} - k\dot{q}]\} \\
& \times \exp\{-\tfrac{1}{2}\nu \int (\beta q^{-2}x^2 - 2\beta^{-1}q\varepsilon + \beta^{-1}k^2 q^2) dt\} \mathcal{D}x \mathcal{D}k \mathcal{D}p \mathcal{D}q \\
= & e^{\nu T/2} \mathcal{N} \int \exp\{-i \int (q + \beta^{-1}q^2\varepsilon)\dot{p} dt\} \\
& \times \exp\{\tfrac{1}{2}\nu \int 2\beta^{-1}q\varepsilon dt\} \exp\{-\tfrac{1}{2\nu} \int [\beta^{-1}q^2\dot{p}^2 + \beta q^{-2}\dot{q}^2] dt\} \mathcal{D}p \mathcal{D}q
\end{aligned}$$

with $N = T/\delta$. The Lie-Trotter product formula was used to go from the second to the third equality. The indices $l + 1/2$ and l indicate that the lattice points do not coincide for x, p or q, k respectively. (This would violate the Heisenberg uncertainty principle.) For the endpoints, the definitions $p_0 := p'$, $p_{N+1} := p''$, $q_0 := q'$ and $q_{N+1} := q''$ were made. Note that $\exp\{\tfrac{1}{2}\nu \delta (-\beta^{-1}\partial_q q^2 \partial_q)\} \exp\{-ik(q-q')\} \approx \exp\{-\tfrac{1}{2}\nu \delta \beta^{-1}k^2 q^2\} \exp\{-ik(q-q')\}$ only to first order in δ , but that was good enough for the path integral. From the second to last to the bottom line, x was substituted by $x + q - \beta^{-1}q^2\varepsilon$ and the x - and p -integrations were carried out.

This formal expression becomes well-defined when understood in the sense of Eq. (88) where the above Wiener measure. Also, the integrals over the Brownian bridges have to be taken in the Stratonovich interpretation of stochastic integrals.

The aim is to show that in the limit of diverging diffusion constant followed by the limit $\varepsilon \rightarrow 0$ the kernel $c_{\beta,\varepsilon}^{-1} K_{\nu,\varepsilon}^0$ reduces to the (weak coherent state) overlap $\langle p''q''|p'q' \rangle$ for $1/4 < \beta \leq 1/2$.

This is similar to what was done in DKP [26]. If one were to follow the proof of DKP [see 2.7] one would have to show these three properties:

- 1) A_ε is a non-negative, self-adjoint operator
- 2) A_ε has an isolated eigenvalue 0
- 3) Pointwise convergence of the kernels $c_{\beta,\varepsilon}^{-1} K_{\nu,\varepsilon}^0$

To copy the proof is not possible. In the “A-world” point 1) could be shown by DKP using Kato-Rellich’s theorem about stability of self-adjointness³⁵. But in the “ A_ε -world” the operator A_ε is not $-\Delta_{LB}$ -bounded. For point 2) and 3) DKP could exploit the connection to the Morse-operator. It is the author’s view that the reason for the existence of this connection between the original 2-dimensional problem on the Lobachevsky half-plane and the 1-dimensional Morse operator lies in the high symmetry of the homogeneous Lobachevsky manifold (constant negative curvature). This connection no longer exists with the ε -modified operator A_ε .

Since other ways for the proof have to be found anyway, it makes sense to reexamine the points 1)-3):

Point 1) certainly is a necessary condition since it ensures non-negativity of A_ε ³⁶ and a real spectrum. It is not obvious why DKP insisted on an isolated eigenvalue 0 unless it was to match the requirements for the application of some theorem which was not mentioned in the article. Of course there must be a discrete ground state with eigenvalue 0 (which exists by construction), but in principle the continuous spectrum could start at 0 as well. In the limit of diverging diffusion constant, the continuous part would disappear. (This was the whole trouble in the preceding attempt to isolate the reproducing kernel for $\beta = 1/2$ by a rescaling procedure.)

Although it is not clear why a gap should be necessary there is every reason to believe that a gap exists. From a physical point of view, the

³⁵If T is self-adjoint, V symmetric and T -bounded with bound smaller than 1, then $T + V$ is self-adjoint.

³⁶See the discussion earlier in this paragraph about operators of the form $B^\dagger B$ which were non-negative only by requirement of certain boundary conditions.

coexistence of a discrete ground state and a continuous state of the same energy would require some conservation law to prevent the discrete state from decaying into the continuum. However, this is invariably associated with a symmetry property of the underlying phase space manifold. Even in the case of the old operator A associated with the Lobachevsky half-plane, a space of high symmetry, there was no sign that anything like that could happen (e.g. in the case $\beta = 1/2$ the spectrum of A started at 0 but was purely continuous; the discrete ground state had disappeared).

So why should one believe that a symmetry which could result in a conservation law could exist in the A_ε -world? One would expect that there is less symmetry than before! From a mathematical point of view one can look at the function $f(\varepsilon)$ the values of which give the starting point of the continuous spectrum. Certainly, $f(0) = (\beta - 1/2)^2$ which is taken from DKP [26]. “No gap” would mean, in this language, that this function f must be discontinuous at 0 and identically 0 in an open neighborhood of 0. It is very hard to imagine how such a functional behavior should arise. If, by any chance, this functional behavior turned up, then it was connected with the special choice of the regularization. But, the regularization is not fixed (the form chosen just seems to be the most simple one) and could e.g. be $\exp\{-q^3\varepsilon^2\}$. This would lead to a different A_ε . Also, the function f would be different. That the new function f should display the same functional behavior as the former one is even more unlikely if not impossible.

To prove the third point DKP used “heavy artillery” (operator estimates based on the connection to the Morse operator) to “shoot a rather small bird”, namely continuity of the limiting function - the reproducing kernel or coherent state overlap³⁷.

Since the connection to the 1-dimensional Morse operator is not available anyway, the natural (and much easier) way is to prove this continuity directly. But, apart from the continuous “factor” $\exp\{-(q+s)\varepsilon\}(qs)^{-\beta}$, the function is analytic in $q^{-1} + i\beta^{-1}p$ and, hence, continuity of the whole function follows.

Summarizing the above, only the self-adjointness of A_ε has to be shown to establish the main result.

The aim is to show that the deficiency index equation $(A_\varepsilon^\dagger\psi)(p, q) = \pm i\psi(p, q)$ has no solution for all $\psi \in D(A_\varepsilon^\dagger)$. This is equivalent to the statement “ A_ε is essentially self-adjoint” [48]. To see this a change of rep-

³⁷There is, of course, a reason why this intricate approach was used. The functional form of $\langle pq|e^{-iT\mathcal{H}}|rs\rangle$ is not known for general Hamiltonian. But, with the operator technique, the pointwise convergence could be shown in general.

representation is made:

$$\psi(p, q) = (2\pi)^{-1/2} \int_{-\infty}^{\infty} e^{ipu} \tilde{\psi}(u, q) du \quad (90)$$

In the new representation the operator A_ε is given by $(A_\varepsilon \phi)(u, q) = \frac{1}{2} \{ \beta(uq^{-1} + 1 + \beta^{-1}q\varepsilon)^2 - 1 - 2\beta^{-1}q\varepsilon - \beta^{-1}\partial_q q^2 \partial_q \} \phi(u, q)$, and the deficiency index equation becomes

$$\{ \beta(uq^{-1} + 1 + \beta^{-1}q\varepsilon)^2 - 1 - 2\beta^{-1}q\varepsilon - \beta^{-1}\partial_q q^2 \partial_q \} \phi(u, q) = \pm 2i \phi(u, q) \quad (91)$$

where $\phi \in D(A_\varepsilon) \Leftrightarrow [\phi \in L^2(\mathbb{R} \times \mathbb{R}^+) \text{ and } (A_\varepsilon \phi) \in L^2(\mathbb{R} \times \mathbb{R}^+)]$.

In the following, u is written as a parameter $[\phi_u(q)]$ to indicate that only q is being dealt with. Multiplying the deficiency index equation from the left by $\phi_u^*(q)$, and integrating over q with integration by parts, one is led to

$$\begin{aligned} & \int \phi_u^*(q) A_\varepsilon \phi_u(q) dq = \pm 2i \int \phi_u^*(q) \phi_u(q) dq \\ \Leftrightarrow & \int \phi_u^*(q) \{ \beta(uq^{-1} + 1 + \beta^{-1}q\varepsilon)^2 - 1 - 2\beta^{-1}q\varepsilon - \beta^{-1}\partial_q q^2 \partial_q \} \phi_u(q) dq \\ & = \pm 2i \int \phi_u^*(q) \phi_u(q) dq \\ \Leftrightarrow & \int [\beta(uq^{-1} + 1 + \beta^{-1}\varepsilon q)^2 - 1 - 2\beta^{-1}\varepsilon q] |\phi_u|^2 dq \\ & + \int \beta^{-1} |\partial_q(q\phi_u)|^2 dq - \beta^{-1} [q|\phi_u|^2]_0^\infty + \beta^{-1} [q^2 \phi_u^* \partial_1 \phi_u]_0^\infty \\ & = \pm 2i \int |\phi_u|^2 dq \end{aligned} \quad (92)$$

In the last equivalence the term on the right hand side is purely imaginary; all the terms on the left hand side of the equality sign are purely real, except maybe for the term $\beta^{-1} [q^2 \phi_u^* \partial_1 \phi_u]_0^\infty$. The goal is to show that this term is either purely real or vanishing. To exhibit this, one has to learn about the asymptotic behavior of $\psi_u(q)$ for extremely large or small values of q .

The Wentzel-Kramers-Brioullin approximation (WKB) can tell about the asymptotic behavior of one-dimensional functions. But, WKB approximately solves $-\partial_x^2 \psi + V(x) - E = 0$ yielding the answer $\psi(x) \approx [V(x) - E]^{-1/4} \exp\{\pm \int^x \sqrt{V(x') - E} dx'\}$ valid in the region to the right of the classically allowed region. The wave function with the positive sign in

the exponent is being discarded as unphysical as it is non-square integrable. The wave solution in the classically allowed region will not be needed. The solution to the left of the classically allowed region is $\psi(x) \approx [V(x) - E]^{-1/4} \exp\{\pm \int_x \sqrt{V(x') - E} dx'\}$ with the positive solution again ignored.

But the problem at hand is not yet in a form to be treated with the standard WKB due to the kinetic energy term $-\beta^{-1} \partial_q q^2 \partial_q$ which stems from the curved Lobachevsky half-plane. One could rather easily develop a new WKB suitable for this problem, but a change of variables, $q \rightarrow a^{-1}$, will bring the problem into standard form. The kinetic energy becomes $-\beta^{-1} a^2 \partial_a^2$ and since $a > 0$ one can divide by $\beta^{-1} a^2$ to obtain the new deficiency equation

$$\begin{aligned} & [\varepsilon^2 a^{-4} + \varepsilon(\beta - 1)a^{-3} + \beta(2\varepsilon u + \beta - 1)a^{-2} + 2\beta^2 u a^{-1} + \beta^2 u^2] \phi_u(a) \\ & - \partial_a^2 \phi_u(a) = \pm 2i a^{-2} \phi_u(a) \end{aligned} \quad (93)$$

For large a ($\hat{=}$ small q), this becomes

$$-\partial_a^2 \phi_u(a) + \beta^2 u^2 \phi_u(a) = 0 \quad (94)$$

with the WKB solution

$$\phi_u(a) \propto \exp\{-\beta|u|a\} \quad (95)$$

For small a ($\hat{=}$ large q) the deficiency index equation reads

$$-\partial_a^2 \phi_u(a) + \varepsilon^2 a^{-4} \phi_u(a) = 0 \quad (96)$$

with WKB solution

$$\phi_u(a) \propto a \exp\{-\varepsilon a^{-1}\} \quad (97)$$

Note how terms introduced by ε dominate in the small a , i.e., large q region, just as one should expect since the regularization of the overlap was aimed at large q . At this point it is already clear that the functions ϕ_u will asymptotically be real, or, to be more precise, they vanish asymptotically. Writing everything in the q -notation again one finds

$$\begin{aligned} \phi_u(q) & \propto \exp\{\beta|u|q^{-1}\} \\ \phi_u(q) & \propto q^{-1} \exp\{-\varepsilon q\} \end{aligned} \quad (98)$$

for the small q and large q asymptotic behavior, respectively. One immediately sees that

$$\begin{aligned} q^2 \phi_u^*(q) \partial_q \phi_u(q) &\propto \exp\{-2\beta|u|q^{-1}\} \xrightarrow{q \rightarrow 0} 0 \\ q^2 \phi_u^*(q) \partial_q \phi_u(q) &\propto \exp\{-2\varepsilon q\} \xrightarrow{q \rightarrow \infty} 0 \end{aligned} \quad (99)$$

showing that the critical term in Eq. (92) indeed vanishes. Thus, the deficiency index equation has no solution. Hence, A_ε is essentially self-adjoint. From now on, and in a not perfectly pure but simpler notation, understand A_ε as the closure of the operator previously called A_ε . The “new” A_ε is self-adjoint. This, together with the fact that A_ε is of the form $B_\varepsilon^\dagger B_\varepsilon$, ensures at the same time that A_ε is non-negative.

Having established these results, one can conclude in the same way DKP did that, for the parameter range $1/4 < \beta \leq 1/2$ and for arbitrary $r \in \mathbb{R}$, $s \in \mathbb{R}^+$,

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} K_{\nu \varepsilon}^0(p'', q'', p', q', T) \\ &= \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} \exp\{-\nu A_\varepsilon T\} \delta(p - p') \delta(q - q')|_{p=p'', q=q''} \\ &= \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} \exp\{\nu T/2\} \int \exp\{-i \int (q + \beta^{-1} q^2 \varepsilon) dp \\ &\quad + \nu \int \beta^{-1} q \varepsilon dt\} d\mu_W^\nu \\ &= \lim_{\varepsilon \rightarrow 0} c_{\beta, \varepsilon}^{-1} \langle p'' q'' | p' q' \rangle_\varepsilon = \langle p'' q'' | p' q' \rangle \end{aligned} \quad (100)$$

The stochastic processes involved are still Brownian bridges, and, when the stochastic integrals are interpreted in the Stratonovich sense, canonical (coordinate) transformations can be made in the same way as before. Thus, the geometric nature of the quantization is untouched. It is just the mathematical structure which is not as elegant as in the case $\beta > 1/2$.

Case $0 < \beta \leq 1/4$

For a parameter $\beta \leq 1/4$, a regularization for large q is not enough. It turns out that an additional p -regularization will even make a regularization for small q necessary (otherwise the overlap would be square integrable, but not in the domain of A_ε).

In the present case, let

$$\langle pq|rs\rangle_\varepsilon := N_\varepsilon \langle pq|rs\rangle e^{-(q+s)\varepsilon - (q^{-1}+s^{-1})\varepsilon - (p^2+r^2)\varepsilon} \quad (101)$$

where $\langle pq|rs\rangle = (qs)^{-\beta} 2^{-2\beta} [(q^{-1}+s^{-1}) + i\beta^{-1}(p-r)]^{-2\beta}$ is the weak coherent state overlap, which is analytic in the complex variable $z := q^{-1} + i\beta^{-1}p$ apart from the factor $(qs)^{-\beta}$. One can write the analytic part (previously called Y - see above) as $e^{(q+s)\varepsilon + (q^{-1}+s^{-1})\varepsilon + (p^2+q^2)\varepsilon} (qs)^\beta \langle pq|rs\rangle_\varepsilon$, and let the differential operator $\partial_{q^{-1}-i\beta^{-1}p} = \frac{1}{2}(-q^2\partial_q + i\beta\partial_p)$ act on this expression. Using $\partial_{z^*}f = 0$ for an analytic function, this results in the new operator

$$B_\varepsilon = \beta^{-1}q\partial_q + 1 + \beta^{-1}q\varepsilon + \beta^{-1}q^{-1}\varepsilon - 2ipq^{-1}\varepsilon - iq^{-1}\partial_p$$

for which $B_\varepsilon \langle pq|rs\rangle_\varepsilon = 0$. As before, define $A_\varepsilon := \frac{1}{2}\beta B_\varepsilon^\dagger B_\varepsilon$, then

$$\begin{aligned} A_\varepsilon = & \frac{1}{2}\{\beta(-iq^{-1}\partial_p + 1 + \beta^{-1}q\varepsilon\beta^{-1}q^{-1}\varepsilon)^2 - 2\beta q^{-2}\varepsilon \\ & + 4ip\partial_q\varepsilon + 4\beta p^2q^{-2}\varepsilon^2 - \beta^{-1}\partial_1 q^2\partial_q - 1 - 2\beta^{-1}q\varepsilon\} \end{aligned} \quad (102)$$

The Feynman-Kac-Stratonovich representation of the kernel of the operator $\exp\{-\nu T A_\varepsilon\}$ is derived in much the same way as before and with the same conventions for notation it reads

$$\begin{aligned} & \exp\{-\nu T A_\varepsilon\} \delta(p-p')\delta(q-q')|_{p=p'', q=q''} \\ = & \exp\left\{-\frac{\nu T}{2}[\beta(-iq^{-1}\partial_p + 1 + \beta^{-1}q\varepsilon\beta^{-1}q^{-1}\varepsilon)^2 - 2\beta q^{-2}\varepsilon \right. \\ & \left. + 4ip\partial_q\varepsilon + 4\beta p^2q^{-2}\varepsilon^2 - \beta^{-1}\partial_1 q^2\partial_q - 1 - 2\beta^{-1}q\varepsilon]\right\} \\ & \times \int \exp\{ix(p-p') - ik(q-q')\} \frac{dxdk}{(2\pi)^2}|_{p=p'', q=q''} \\ = & e^{\nu T/2} \lim_{N \rightarrow \infty} [\exp\{-\nu\delta/2 \\ & [\beta(-iq^{-1}\partial_p + 1 + \beta^{-1}q\varepsilon + \beta^{-1}q^{-1}\varepsilon)^2 - 2\beta q^{-2}\varepsilon - 2\beta^{-1}q\varepsilon]\} \\ & \times \exp\{-\nu\delta/2(-\beta^{-1}\partial_q q^2\partial_q)\} \exp\{-\nu\delta/2 \cdot 4ip\partial_q\varepsilon\} \\ & \exp\{-\nu\delta/2 \cdot 4\beta p^2q^{-2}\varepsilon^2\}]^N \\ & \int \exp\{ix(p-p') - ik(q-q')\} \frac{dxdk}{(2\pi)^2}|_{p=p'', q=q''} \end{aligned}$$

$$\begin{aligned}
&= \lim_{N \rightarrow \infty} e^{\nu T/2} \int \exp\{i \sum x_{l+1/2}(p_{l+1} - p_l) - ik_{l+1/2}(q_{l+1} - q_l)\} \\
&\quad \times \exp\{-\nu\delta/2 \sum [\beta(q_l^{-1}x_{l+1/2} + 1 + \beta^{-1}q_l\varepsilon + \beta^{-1}q_l^{-1}\varepsilon)^2 \\
&\quad - 2\beta q_l^{-2}\varepsilon - 2\beta^{-1}q_l\varepsilon]\} \\
&\quad \times \exp\{-\nu\delta/2 \sum \beta^{-1}k_{l+1/2}^2 q_l^2\} \exp\{-\nu\delta/2 \sum 4ip_l(-ik_{l+1/2})\varepsilon\} \\
&\quad \exp\{-\nu\delta/2 \sum 4\beta p_l^2 q_l^{-2}\varepsilon^2\} \prod_{l=0}^N \frac{dk_{l+1/2} dx_{l+1/2}}{(2\pi)^2} \prod_{l=1}^N dp_l dq_l \\
&=: e^{\nu T/2} \mathcal{N} \int \exp\{i \int (x\dot{p} - k\dot{q})dt\} \\
&\quad \times \exp\{-\nu/2 \int \{\beta(q^{-1}x + 1 + \beta^{-1}q\varepsilon + \beta^{-1}q^{-1}\varepsilon)^2 - 2\beta q^{-2}\varepsilon - 2\beta^{-1}q\varepsilon \\
&\quad + \beta^{-1}k^2 q^2 + 4pk\varepsilon + 4\beta p^2 q^{-2}\varepsilon^2\}dt\} \mathcal{D}x \mathcal{D}k \mathcal{D}p \mathcal{D}q \\
&= e^{\nu T/2} \mathcal{N} \int \exp\{i \int [(x - q - \beta^{-1}q^2\varepsilon - \beta^{-1}\varepsilon)\dot{p} - k\dot{q}]dt\} \\
&\quad \times \exp\{-\nu/2 \int (\beta q^{-2}x^2 - 2\beta q^{-2}\varepsilon - 2\beta^{-1}q\varepsilon + \beta^{-1}k^2 q^2 + 4pk\varepsilon \\
&\quad + 4\beta p^2 q^{-2}\varepsilon^2)dt\} \mathcal{D}x \mathcal{D}k \mathcal{D}p \mathcal{D}q \\
&= e^{\nu T/2} \mathcal{N} \int \exp\{-i \int (q + \beta^{-1}\varepsilon + \beta^{-1}q^2\varepsilon)\dot{p}dt + i \int 2\beta pq^{-2}\dot{q}dt\} \\
&\quad \times \exp\{\nu/2 \int (2\beta^{-1}q\varepsilon + 2\beta q^{-2}\varepsilon - 4\beta p^2 q^{-2}\varepsilon^2)dt\} \\
&\quad \exp\{-1/(2\nu) \int [\beta^{-1}q^2\dot{p}^2 + \beta q^{-2}\dot{q}^2]dt\} \mathcal{D}p \mathcal{D}q
\end{aligned}$$

Introducing the Wiener measure, as above [see Eq. (89)], and performing a partial integration on $2\beta\varepsilon pq^{-2}dq = -2\beta\varepsilon pd(q^{-1}) = -2\beta\varepsilon pq^{-1}|_{(p',q')}^{(p'',q'')} + 2\beta\varepsilon q^{-1}dp$, the final result is

$$\begin{aligned}
&K_{\nu,\varepsilon}^0(p'', p', q'', q') \\
&:= \exp\{\nu T/2\} \exp\{-i\beta^{-1}\varepsilon(p'' - p') - 2i\beta\varepsilon(p''q''^{-1} - p'q'^{-1})\} \\
&\quad \times \int \exp\{-i \int (q + \beta^{-1}q^2\varepsilon - 2\beta q^{-1}\varepsilon)dp \\
&\quad + \nu \int [\beta q^{-2}\varepsilon + \beta^{-1}q\varepsilon - 2\beta p^2 q^{-2}\varepsilon^2]dt\} d\mu_W^\nu
\end{aligned} \tag{103}$$

As before, the aim is to establish $\lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta,\varepsilon}^{-1} K_{\nu,\varepsilon}^0(p'', p', q'', q') = \langle p''q''|p'q' \rangle$. The questions about “gap/no gap” and about pointwise con-

vergence can be answered in the same way as in the previous section. The self-adjointness could previously be proved since the “old” A_ε did not contain p explicitly. Instead of trying to solve the deficiency index equation for the “new” A_ε one can avoid this question altogether.

If A_ε is self-adjoint there is nothing to show. Assume A_ε is not self-adjoint. The (sesquilinear) form $s_\varepsilon(x, y) := \langle x | A_\varepsilon y \rangle$, generated by A_ε , is closable since A_ε is symmetric and bounded below [48]. There is a bijection between the set of all (densely defined) closed, below-bounded forms and the set of all self-adjoint, below-bounded operators. Let $\bar{s}_\varepsilon$ be the closure of the form generated by A_ε and $A_{\bar{s}_\varepsilon}$ be the self-adjoint operator associated with $\bar{s}_\varepsilon$. Then $A_{\bar{s}_\varepsilon}$ preserves the lower bound and is called the Friedrichs’ extension of the operator A_ε . It is the unique extension fulfilling $D(A_{\bar{s}_\varepsilon}) \subset D(\bar{s}_\varepsilon)$ [48].

In a slight abuse of notation, $A_{\bar{s}_\varepsilon}$ will be written as A_ε . So from now on A_ε denotes the Friedrichs’ extension (which is trivial in the case that A_ε is already self-adjoint). Then it is clear that A_ε is non negative.

With this in mind, one can establish the desired result without further work. The phase factors in Eq. (103) are ν -independent, so they come outside of the ν -limit and become unity.

$$\begin{aligned}
& \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} K_{\nu, \varepsilon}^0(p'', p', q'', q') \\
&:= \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} \exp\{\nu T/2\} \int \exp\{-i \int (q + \beta^{-1} q^2 \varepsilon - 2\beta q^{-1} \varepsilon) dp \\
&\quad + \nu \int [\beta q^{-2} \varepsilon + \beta^{-1} q \varepsilon - 2\beta p^2 q^{-2} \varepsilon^2] dt\} d\mu_W^\nu \\
&= \langle p'' q'' | p' q' \rangle
\end{aligned} \tag{104}$$

This is a valid path integral representation for all $0 < \beta \leq 1/4$.

3.2.4 Introducing dynamics

Dynamics are introduced in the same way as for the spectral approach. For a linear Hamiltonian $\mathcal{H} = RQ + SD$, the problem is already solved since it reduces to an overlap with modified ending points. What remains to do is to write down the path integral. This is straightforward since everything stated previously concerning the measure, etc., remains valid, and the formula for $1/4 < \beta \leq 1/2$ is

$$\begin{aligned}
& \langle p'' q'' | e^{-i(RQ+SD)T} | p' q' \rangle = \langle p'' e^{ST} + \frac{R}{S}(e^{ST} - 1), q'' e^{-ST} | p' q' \rangle \\
& = \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} e^{\nu T/2} \int_{p', q'}^{p'', q''} \exp\{-i \int (q + \beta^{-1} q^2 \varepsilon) dp \\
& \quad + \nu \int \beta^{-1} q \varepsilon dt\} d\mu_W^{\nu} \\
& = \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} e^{\nu T/2} \int_{p', q'}^{p'', q''} \exp\{-i \int (q e^{-St} + \beta^{-1} q^2 e^{-2St} \varepsilon) \\
& \quad \times d[pe^{St} + \frac{R}{S}(e^{St} - 1)] + \nu \int \beta^{-1} q e^{-St} \varepsilon dt\} d\mu_W^{\nu} \\
& = \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} e^{\nu T/2} \int_{p', q'}^{p'', q''} \exp\{-i [\int (q + \beta^{-1} q^2 e^{-St} \varepsilon) dp \\
& \quad + \int (q + \beta^{-1} q^2 e^{-St} \varepsilon)(Sp + R) dt] + \nu \int \beta^{-1} q e^{-St} \varepsilon dt\} d\mu_W^{\nu} \quad (105)
\end{aligned}$$

The complexity of this expression can be hidden by introducing the variable $q_\varepsilon := q + \beta^{-1} q^2 e^{-St} \varepsilon$ and the measure $d\mu_W^{\nu, \varepsilon} := \exp\{\nu \int \beta^{-1} q e^{-St} \varepsilon dt\} d\mu_W^{\nu}$. The new measure is equivalent to the old Wiener measure since the factor $\exp\{\nu \int \beta^{-1} q e^{-St} \varepsilon dt\}$ is a Radon-Nykodym derivative. Then, the formula resembles the path integral for coherent states:

$$\lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} e^{\nu T/2} \int_{p', q'}^{p'', q''} \exp\{-i [\int q_\varepsilon dp + (Spq_\varepsilon + Rq_\varepsilon) dt] d\mu_W^{\nu, \varepsilon} \quad (106)$$

The ε -modified Hamiltonian is given by the weak modified symbol $h_{w, \varepsilon} := Rq_\varepsilon + Spq_\varepsilon$.

The same procedure for $0 < \beta \leq 1/4$ leads to:

$$\begin{aligned}
& \langle p'' q'' | e^{-i(RQ+SD)T} | p' q' \rangle \\
& = \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} e^{\nu T/2} \int_{p', q'}^{p'', q''} \exp\{-i [\int (q + \beta^{-1} q^2 e^{-St} \varepsilon - 2\beta q^{-1} e^{2St}) dp \\
& \quad + \int (q + \beta^{-1} q^2 e^{-St} \varepsilon - 2\beta q^{-1} e^{2St})(Sp + R) dt] \\
& \quad + \nu \int (\beta^{-1} q e^{-St} \varepsilon + \beta q^{-2} e^{2St} \varepsilon - 2\beta p^2 q^{-2} e^{4St} \varepsilon^2) dt\} d\mu_W^{\nu} \\
& = \lim_{\varepsilon \rightarrow 0} \lim_{\nu \rightarrow \infty} c_{\beta, \varepsilon}^{-1} e^{\nu T/2} \int_{p', q'}^{p'', q''} \exp\{-i [\int \tilde{q}_\varepsilon dp + \int (Sp\tilde{q}_\varepsilon + R\tilde{q}_\varepsilon) dt] d\tilde{\mu}_W^{\nu, \varepsilon} \quad (107)
\end{aligned}$$

The variable $\tilde{q}_\varepsilon := (q + \beta^{-1}q^2e^{-St}\varepsilon - 2\beta q^{-1}e^{2St})$ and the Radon-Nykodym measure $d\tilde{\mu}_W^{\nu,\varepsilon} := \exp\{\nu \int (\beta^{-1}qe^{-St}\varepsilon + \beta q^{-2}e^{2St}\varepsilon - 2\beta p^2q^{-2}e^{4St}\varepsilon^2)dt\} d\mu_W^\nu$ were used. The weak modified symbol is now $h_{w,\varepsilon} = R\tilde{q}_\varepsilon + Sp\tilde{q}_\varepsilon$.

The question, how this can be extended to, say, all polynomial Hamiltonians, was already discussed in subsection (3.2.2). The remaining questions are the same.

4 Final Remarks

In this work, a path integral with Wiener measure for weak coherent states was constructed, but some questions still remain.

The path integral with Wiener measure for the coherent states basically contained two “ingredients”, both of which were of geometric nature. The canonical one-form, $\theta = p dq$, (where $p dq$ is locally valid if θ is expressed in canonical coordinates) as a phase in the integrand, and a (Riemannian) metric, $d\sigma^2$, in a regularization factor, by virtue of which the Wiener measure emerged.

These appealing features are maintained in the spectral approach to a weak coherent state path integral, but, unfortunately, it turned out that this approach could be applied to one special case only. The regularized approach on the other hand introduced many new terms, and spoils the beauty of the original idea. So the question is whether one can avoid this. Naturally, it is possible to try different regularizations. One way would be a cut-off method. Instead of the normalization condition (which was restored by the ε -modification), one could confine the problem to the square $[-L, L]^2$ and impose appropriate boundary conditions³⁸. The functions are periodically continued outside of the square $[-L, L]^2$, while the boundary conditions are enforced by sums of δ -functions. The limit $L \rightarrow \infty$ is now the final step of the whole procedure. This has not been worked out yet, but conceptual studies on one-dimensional problems indicate that the final path integral seems to be closer to the old coherent state path integral with Wiener measure (at least under esthetic aspects).

Another question, already raised, is the maximal class of Hamiltonians for which the affine weak coherent state path integral is valid. It was rigorously established only for linear Hamiltonians. Its validity for Hamiltonians polynomial in the kinematical operators Q and D was only conjectured. However, for the regularized approach there could be another way to prove this. This proof was used by Daubechies and Klauder [39] and by DKP [26] in the coherent state case for $\beta > 1/2$ and basically involves the same steps as in the case of zero Hamiltonian. The operator νA_ε has to be replaced by $\nu A_\varepsilon + i h_{\nu arepsilon}$. The symbol $h_{\nu arepsilon}$, representing the Hamiltonian, could be different from the weak modified symbol introduced in the last section. But, a rigorous treatment of this problem is yet to be done.

Finally, the question can be addressed as to whether the weak coherent state path integral with Wiener measure is amenable for computational

³⁸The non-negative, affine variable is supposed to be mapped to the whole real line first.

pursuits. Calculations using Monte Carlo methods have been done for the coherent state path integral with Wiener measure [49], but the results were not too promising due to the slow convergence encountered. The same can be assumed for the weak coherent state path integral, which is already more complicated in its analytic form. So it might turn out that the merits of this path integral are rather confined to the theoretical aspect.

A Appendix

A.1 The affine group

Since the affine group is central to the whole present work more of its properties are listed here for the interested reader.

The affine group is a subgroup of $SU(1, 1)$ [28]. To see this, recall first that the generators Q (with $Q > 0$) and D fulfill the affine commutation relation $[Q, D] = iQ$. Define the operator

$$R := DQ^{-1}D + [(\beta - \frac{1}{2})^2 - \frac{1}{4}]Q^{-1}$$

and the operators

$$\begin{aligned} K_0 &:= \frac{1}{2}(\beta Q + \beta^{-1}R) \\ K_1 &:= \frac{1}{2}(\beta Q - \beta^{-1}R) \\ K_2 &:= -D \end{aligned}$$

Using $[Q^\alpha, D] = i\alpha Q^\alpha$, it follows that $[R, D] = -iR$, and further

$$\begin{aligned} [K_1, K_2] &= -iK_0 \\ [K_0, K_1] &= iK_2 \\ [K_2, K_0] &= iK_1 \end{aligned}$$

This is the Lie algebra of $SU(1, 1)$. Hence, $Q = \beta^{-1}(K_0 + K_1)$ and $D = -K_2$ form a subalgebra, and the affine group is a subgroup of $SU(1, 1)$.

For the family $|\eta_\beta\rangle$ of minimum uncertainty states introduced in section (2.3), one has

$$\begin{aligned} K_0|\eta_\beta\rangle &= \beta|\eta_\beta\rangle \\ K_+|\eta_\beta\rangle &= 0 \end{aligned}$$

where $K_\pm := K_2 \pm K_1$. These properties can most easily be checked in x -representation in which $R = -\partial_x x \partial_x + (\beta - 1/2)^2 x^{-1}$.

For the resolution of unity, the left-invariant group measure is important. And the fact that the affine group can generate weak coherent states (for certain choices of the fiducial vector) is due to the non-equality of the left- and right-invariant group measure.

In general, the left-invariant group measure can be determined as follows: Let $U(g)$ be a group representation, X_a , $a \in \{1, \dots, A\}$ the group generators, and g_0 some group element, then $U^{-1}(g)dU(g) = X_a\omega^a(g) = X_a\omega^a(g_0g)$, where the ω^a are one-forms. The left invariant group measure is (proportional to) $d\mu_L(g) = \omega^1(g) \wedge \dots \wedge \omega^A(g) = d\mu_L(g_0g)$.

For the right-invariant group measure, one has (with the same notation): $[dU(g)]U^{-1}(g) = X_a\Omega^a(g) = X_a\Omega^a(gg_0)$, where the Ω^a are again one-forms, and the right-invariant group measure is (proportional to) $d\mu_R(g) = \Omega^1(g) \wedge \dots \wedge \Omega^A(g) = d\mu_R(gg_0)$.

Using the group representation $U(p, q) = e^{ipQ}e^{-i\ln qD}$ of the affine group [which is familiar from section (2.3)], one finds for the left-invariant group measure $U^{-1}dU = i(Qqdp - Dq^{-1}dq)$, and, hence,

$$d\mu_L(p, q) = qdp \wedge q^{-1}dq = dpdq$$

For the right-invariant group measure, one has $(dU)U^{-1} = i[Q(dp + pq^{-1}dq) - Dq^{-1}dq]$, and, hence,

$$d\mu_R(p, q) = (dp + pq^{-1}dq) \wedge q^{-1}dq = q^{-1}dpdq$$

This is not equal to the left-invariant group measure.

A.2 Parameters in DKP and the present work

As mentioned before, the parameter β is different in its physical meaning in the present work compared to the DKP-article [26], but mathematically it turns out to be equivalent. To make comparison easier the connection will be studied in the following.

Start with the original 2-parameter family of minimum uncertainty states for the kinematical variables D and Q

$$\eta_{\alpha\beta'}(x) = N_{\alpha\beta'}x^\alpha e^{-\beta'x}$$

The norm of these states can be computed:

$$\begin{aligned} 1 &\stackrel{!}{=} \langle \eta | \eta \rangle = \int_0^\infty dx N_{\alpha\beta'}^2 x^{2\alpha} e^{-2\beta'x} \\ &= N_{\alpha\beta'}^2 (2\beta')^{-2\alpha-1} \int_0^\infty dx x^{2\alpha} e^{-x} = N_{\alpha\beta'}^2 (2\beta')^{-2\alpha-1} \Gamma(2\alpha + 1) \end{aligned}$$

W.l.o.g., choose $N_{\alpha\beta'}$ real. So $N_{\alpha\beta'}^2 = (2\beta')^{2\alpha+1} \Gamma^{-1}(2\alpha + 1)$, and a similar calculation shows $\langle Q \rangle = \langle \eta | Q | \eta \rangle = N_{\alpha\beta'}^2 (2\beta')^{2\alpha-2} \Gamma(2\alpha + 2) = (\beta')^{-1}(\alpha +$

1/2). Setting $\langle Q \rangle = k$ for the mean of Q , this means $\alpha = \beta'k - 1/2$ and one is left with a family of minimum uncertainty states:

$$\eta_{\beta'k}(x) = N_{\beta'k} x^{\beta'k-1/2} e^{-\beta'x}$$

The choice in this work is $k = 1$ and $\beta := \beta'$, whereas DKP [26] chose $k = \beta$ and $\beta' = 1$.

Let $U(p, q) = e^{ipQ} e^{-i \ln q D}$ and $|pq\rangle = U(p, q)|\eta\rangle$ (DKP: $p \rightarrow b, q \rightarrow a^{-1}$), then the coherent state overlap becomes:

$$\begin{aligned} \langle pq|rs\rangle &= \langle \eta | e^{i \ln q D} e^{-ipQ} | e^{irQ} e^{-i \ln s D} | \eta \rangle \\ &= \int_0^\infty dx [e^{ipx} e^{-i \ln q (-ix\partial_x - i/2)} \eta]^* [e^{irx} e^{-i \ln s (-ix\partial_x - i/2)} \eta(x)] \\ &= (qs)^{-1/2} \int_0^\infty dx \eta^*(x/q) e^{-ix(p-r)} \eta(x/s) \\ &= (qs)^{-1/2} N_{\beta'k}^2 q^{1/2-\beta'k} s^{1/2-\beta'k} \\ &\quad \times \int_0^\infty dx x^{2\beta'k-1} e^{-\beta'x[(q^{-1}+s^{-1})+i\beta'^{-1}(p-r)]} \\ &= (qs)^{-\beta'k} (2\beta')^{2\beta'k} \Gamma^{-1}(2\beta'k) (\beta')^{-2\beta'k} \\ &\quad \times [(q^{-1}+s^{-1})+i\beta'^{-1}(p-r)]^{-2\beta'k} \int_0^\infty dx x^{2\beta'k-1} e^{-x} \\ &= (qs)^{-\beta'k} 2^{2\beta'k} [(q^{-1}+s^{-1})+i\beta'^{-1}(p-r)]^{-2\beta'k} \end{aligned}$$

[Here: $\langle pq|rs\rangle = (qs)^{-\beta} 2^{2\beta} [(q^{-1}+s^{-1})+i\beta^{-1}(p-r)]^{-2\beta}$

DKP: $\langle ab|a'b'\rangle = (aa')^\beta 2^{2\beta} [(a+a')+i(b-b')]^{-2\beta}$]

Define the operator

$$B := -ik^{-1}q^{-1}\partial_p + 1 + (k\beta')^{-1}q\partial_q$$

[Here: $B = -iq^{-1}\partial_p + 1 + \beta^{-1}q\partial_q$

DKP: Note: $\partial_q \rightarrow -a^2\partial_a = \partial_{a^{-1}}$, so $B = -i\beta^{-1}a\partial_b + 1 - \beta^{-1}a\partial_a$]

Direct computation shows $B\langle pq|rs\rangle = 0$. By direct calculation [e.g. in x -representation, where $Q = x$ and $D = -i(x\partial_x + \frac{1}{2})$] one can easily verify

$$(\frac{Q}{k} - 1 + \frac{iD}{k\beta'})|\eta\rangle = 0$$

and so one gets even for arbitrary ψ (with the use of $e^{i \ln q D} Q e^{-i \ln q D} = qQ$):

$$B\langle pq|\psi\rangle = B\langle \eta | e^{i \ln q D} e^{-ipQ} |\psi\rangle = \langle \eta | [\frac{i}{\beta'k} D + 1 - \frac{1}{k} Q] e^{i \ln q D} e^{-ipQ} |\psi\rangle = 0$$

DKP chose a different factor $c = (k\beta')^2 = \beta^2$ for the operator $A = c \cdot B^\dagger B$ than was selected for this work ($c = \frac{1}{2}k\beta' = \frac{1}{2}\beta$).

So here [compare Eq. (50)]:

$$A = \frac{1}{2}\{-\beta^{-1}\partial_q q^2 \partial_q - \beta q^{-2} \partial_p^2 - 2i\beta q^{-1} \partial_p + \beta - 1\}$$

and DKP has (with their a - b notation):

$$A = -a^2(\partial_a^2 + \partial_b^2) - 2i\beta a \partial_b + \beta^2 - \beta$$

A.3 Spectrum of A

Case $1/4 < \beta \leq 1/2$

To show that there is no sequence ψ_n which fulfills the necessary requirements, one starts with the general function

$$\psi_n(q) = N_n [1 + \sum_{m=1}^M a_m q^{\sigma_m} / n^{\rho_m}] e^{-\sum_{k=1}^K c_k q^{\mu_k} / n^{\nu_k}} \psi(q)$$

In principle $(\rho_m)_n$ etc. could be sequences themselves (which have to fulfill some requirements, e.g. $n^{\rho_m(n)}$ must go to infinity), but since it does not matter for the later calculations it is enough to think about them as constant sequences (or constants). Then one must have $\rho_m, \nu_k, c_k \in \mathbb{R}^+$, and $\mu_k, a_m \in \mathbb{R}$ for all m or k respectively. If there is a k such that $\mu_k < 0$ (regularization for small q), then $\sigma_m \in \mathbb{R}^{\mathbb{N}}$. Else, there is a constant $c(\beta)$, depending on β , which must ensure $\psi_n, A\psi_n \in L^2$ when setting $\sigma_m \geq c(\beta)$.

The function ψ_n is square integrable and normalized with normalization constant N_n , and ψ_n/N_n converges to $\psi(q)$ in the limit $n \rightarrow \infty$. If one can show that $\|A\psi_n\| \not\rightarrow 0$ for $n \rightarrow \infty$, then one can conclude by a density argument that there is no function which will do the job.

Define $1 + \sum_m := 1 + \sum_{m=1}^M a_m q^{\sigma_m} / n^{\rho_m}$, $\sum'_m := \sum_{m=1}^M a_m \sigma_m q^{\sigma_m-1} / n^{\rho_m}$, $\sum''_m := \sum_{m=1}^M a_m \sigma_m (\sigma_m - 1) q^{\sigma_m-2} / n^{\rho_m}$, $\sum_k := \sum_{k=1}^K c_k q^{\mu_k} / n^{\nu_k}$, $\sum'_k := \sum_{k=1}^K c_k \mu_k q^{\mu_k-1} / n^{\nu_k}$ and $\sum''_k := \sum_{k=1}^K c_k \mu_k (\mu_k - 1) q^{\mu_k-2} / n^{\nu_k}$ then:

$$\begin{aligned}
& A\psi_n \\
= & \{-q^2(\frac{\sum'_m}{1+\sum_m})^2 - q^2(\sum'_k)^2 + 2\beta q(\frac{\sum'_m}{1+\sum_m}) - 2\beta q(\sum'_k) \\
& -4\beta[+]^{-1}(\frac{\sum'_m}{1+\sum_m}) + 4\beta[+]^{-1}(\sum'_k) + 2q^2(\frac{\sum'_m}{1+\sum_m})(\sum'_k) \\
& -q^2\frac{(1+\sum_m)(\sum''_m) - (\sum'_M)^2}{(1+\sum_m)^2} + q^2(\sum''_k) - 2q(\frac{\sum'_m}{1+\sum_m}) + 2q(\sum'_k)\}\psi_n
\end{aligned}$$

Since the norm $\|A\psi_n\|$ has to be calculated one needs:

$$\begin{aligned}
& (A\psi_n)^*(A\psi_n) \\
= & [\{-q^2(\frac{\sum'_m}{1+\sum_m})^2 - q^2(\sum'_k)^2 + 2\beta q(\frac{\sum'_m}{1+\sum_m}) - 2\beta q(\sum'_k) \\
& + 2q^2(\frac{\sum'_m}{1+\sum_m})(\sum'_k) - q^2\frac{(1+\sum_m)(\sum''_m) - (\sum'_M)^2}{(1+\sum_m)^2} \\
& + q^2(\sum''_k) - 2q(\frac{\sum'_m}{1+\sum_m}) + 2q(\sum'_k)\}^2 \\
& - \{\dots\}4\beta(\frac{\sum'_m}{1+\sum_m})\frac{[+] + [-]}{[]} + \{\dots\}4\beta(\sum'_k)\frac{[+] + [-]}{[]} \\
& 16\beta^2[]^{-1}(\frac{\sum'_m}{1+\sum_m})^2 + 16\beta^2[]^{-1}(\sum'_k)^2 - 32\beta^2[]^{-1}(\frac{\sum'_m}{1+\sum_m})(\sum'_k)] \\
& \times \psi_n^* \psi_n
\end{aligned}$$

where $\{\dots\}$ was used as an abbreviation for the terms in braces in the lines 2 to 4.

Next, proceed by calculating the norm of ψ_n :

$$\begin{aligned}
& \|\psi_n\|^2 \\
= & N_n^* N_n \cdot \text{const} \int_0^\infty dq (1+\sum_m)^2 e^{-2(\sum_k)} q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \\
& \times \int_{-\infty}^\infty dp [1 + \beta^{-2} p^2]^{-2\beta} \\
= & \text{const} \cdot N_n^* N_n [\int_0^c dq (1+\sum_m)^2 e^{-2(\sum_k)} q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \\
& + \int_c^\infty dq (1+\sum_m)^2 e^{-2(\sum_k)} q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta}]
\end{aligned}$$

In the third line distinguish two cases: i) There is no $\mu_k < 0$, i.e., there is no small q regularization and restrictions to (σ_m) apply. Then set $c = 0$, i.e., the first integral vanishes. ii) If there exists a $\mu_k < 0$, then choose the biggest μ_k/ν_k which is smaller than zero and call it μ_{max}/n_{max} . In the fourth line choose the smallest μ_k/ν_k which is bigger than zero and call it μ_{min}/n_{min} . Then the equation continues:

$$\begin{aligned}
& \dots \\
& = \text{const} \cdot N_n^* N_n \left\{ \int_0^c dq (1 + \sum_m)^2 \exp\left[-\frac{2c_{max}}{q^{|\mu_{max}|} n^{\nu_{max}}}\right] e^{-2(\sum_{k \neq k_{max}})} \right. \\
& \quad \times q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \\
& \quad + \int_c^\infty dq (1 + \sum_m)^2 \exp\left[-\frac{2c_{min} q^{\mu_{min}}}{n^{\nu_{min}}}\right] e^{-2(\sum_{k \neq k_{min}})} \\
& \quad \times q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} \Big\} \\
& = \text{const} \cdot N_n^* N_n \left\{ \int_0^c dq \left(1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{-\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m}\right)^2 \right. \\
& \quad \times \exp\left[-\frac{2c_{max}}{q^{|\mu_{max}|}}\right] \exp\left[-2 \sum_{k \neq k_{max}} c_k q^{\mu_k} n^{-\nu_{max}/|\mu_{max}| \cdot \mu_k - \nu_k}\right] \\
& \quad \times q^{-2\beta} n^{(\nu_{max}/|\mu_{max}|)(2\beta-1)} (n^{\nu_{max}/|\mu_{max}|} q^{-1} + s^{-1})^{1-4\beta} \\
& \quad + \int_c^\infty dq \left(1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{\nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m}\right)^2 \\
& \quad \times \exp\left[-2c_{min} q^{\mu_{min}}\right] \exp\left[-2 \sum_{k \neq k_{min}} c_k q^{\mu_k} n^{\nu_{min}/\mu_{min} \cdot \mu_k - \nu_k}\right] \\
& \quad \times q^{-2\beta} n^{-(\nu_{min}/\mu_{min})(2\beta-1)} [(n^{\nu_{min}/\mu_{min}} q)^{-1} + s^{-1}]^{1-4\beta} \Big\} \\
& \leq \text{const} \cdot N_n^* N_n \left\{ \text{const} \cdot n^{-\nu_{max}/|\mu_{max}| \cdot 2\beta} \right. \\
& \quad \times \int_0^c dq \left(1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{-\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m}\right)^2 \exp\left[-\frac{2c_{max}}{q^{|\mu_{max}|}}\right] q^{2\beta-1} \\
& \quad + \text{const} \cdot n^{(\nu_{min}/\mu_{min})(1-2\beta)} \\
& \quad \times \int_c^\infty dq \left(1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{\nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m}\right)^2 \exp\left[-2c_{min} q^{\mu_{min}}\right] q^{-2\beta} \Big\}
\end{aligned}$$

By choosing $\mu_{max}, \nu_{max}, \mu_{min}$ and ν_{min} as above, one ensures that after a substitution ($q \rightarrow q n^{-\nu_{max}/|\mu_{max}|}$ for small q and $q \rightarrow q n^{\nu_{min}/\mu_{min}}$ for large q), the remaining summands of the k -sum quickly go to zero, i.e., their

exponentials go to 1 from below. Hence, they can be approximated by one. Observe that the different q -dependence ($q^{-2\beta}$ in the last and $q^{2\beta-1}$ in the second to last line) in the integrals stems from the factor $(q^{-1} + s^{-1})^{1-4\beta}$. After the different substitutions in the two regions, this leads to different integrals and to different n -dependences in front of the integrals as well.

So,

$$\begin{aligned}
& N_n^* N_n \\
\geq & \{const \cdot n^{-\nu_{max}/|\mu_{max}| \cdot 2\beta} \\
& \times \int_0^c dq (1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{-\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m})^2 \exp[-\frac{2c_{max}}{q^{|\mu_{max}|}}] q^{2\beta-1} \\
& + const \cdot n^{(\nu_{min}/\mu_{min})(1-2\beta)} \\
& \times \int_c^\infty dq (1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{\nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m})^2 \exp[-2c_{min} q^{\mu_{min}}] q^{-2\beta}\}^{-1}
\end{aligned}$$

and this is the correct asymptotic behavior.

Consider $(A\psi_n)^*(A\psi_n)$ again. This expression naturally is ≥ 0 . Certainly, $A\psi_n$ can be 0 for some n (even infinitely many, as long as there are infinitely many for which this is not true), and one chooses w.l.o.g. the subsequence for which $A\psi_n \neq 0$ and refers to it by the same name (i.e. the sequence will still be labeled by n). Then $\|A\psi_n\| > 0$ for all n and the goal is to show that $\lim_{n \rightarrow \infty} \|A\psi_n\| \neq 0$.

Pick the summand $q^2(\sum'_k)^2$ of $(A\psi_n)^*(A\psi_n)$. The p -integration will give $const \cdot (q^{-1} + s^{-1})^{1-4\beta}$. So compute the expression:

$$N_n^* N_n \int_0^\infty dq q^{-2\beta} (q^{-1} + s^{-1})^{1-4\beta} (1 + \sum_m)^2 e^{-2(\sum_k)} q^2 (\sum'_k)^2$$

$$\begin{aligned}
&\geq \left\{ \int_0^c dq n^{-\nu_{max}/|\mu_{max}| \cdot (2\beta-1)} q^{-2\beta} (n^{\nu_{max}/|\mu_{max}|} q^{-1} + s^{-1})^{1-4\beta} \right. \\
&\quad \times \exp\left[-\frac{2c_{max}}{q^{|\mu_{max}|}}\right] e^{-2(\sum_{k \neq k_{max}})} \left(\sum_k c_k \mu_k q^{\mu_k} n^{-\nu_{max}/|\mu_{max}| \cdot \mu_k - \nu_k}\right)^2 \\
&\quad \times \left(1 + \sum_m a_m q^{\sigma_m} n^{-\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m}\right)^2 \\
&\quad + \int_c^\infty dq n^{\nu_{min}/\mu_{min} \cdot (1-2\beta)} [(qn^{\nu_{min}/\mu_{min}})^{-1} + s^{-1}]^{1-4\beta} \\
&\quad \times \exp[-2c_{min} q^{\mu_{min}}] e^{-2(\sum_{k \neq k_{min}})} \left(\sum_k c_k \mu_k q^{\mu_k} n^{\nu_{min}/\mu_{min} \cdot \mu_k - \nu_k}\right)^2 \\
&\quad \times \left(1 + \sum_m a_m q^{\sigma_m} n^{\nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m}\right)^2 \Big\} \\
&\quad \times \{const \cdot n^{-\nu_{max}/|\mu_{max}| \cdot 2\beta} \\
&\quad \times \int_0^c dq \left(1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{-\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m}\right)^2 \exp\left[-\frac{2c_{max}}{q^{|\mu_{max}|}}\right] q^{2\beta-1} \\
&\quad + const \cdot n^{(\nu_{min}/\mu_{min})(1-2\beta)} \\
&\quad \times \int_c^\infty dq \left(1 + \sum_{m=1}^M a_m q^{\sigma_m} n^{\nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m}\right)^2 \exp[-2c_{min} q^{\mu_{min}}] q^{-2\beta} \Big\}^{-1}
\end{aligned}$$

Define ρ_{min}, σ_{min} by: $-\nu_{max}/|\mu_{max}| \cdot \sigma_{min} - \rho_{min} \geq -\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m$ for all m and define ρ_{max}, σ_{max} by: $\nu_{min}/\mu_{min} \cdot \sigma_{max} - \rho_{max} \geq \nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m$ for all m.

The n -dependence is discussed with the help of these definitions:

1.term $[\int_0^c / (\int_0^c + \int_c^\infty)]$ contains:

Numerator:

- a) $n^{-\nu_{max}/|\mu_{max}| \cdot 2\beta}$
- b) the “worst” term of $(\sum_k c_k \mu_k q^{\mu_k} n^{-\nu_{max}/|\mu_{max}| \cdot \mu_k - \nu_k})^2$ is the one where $k = k_{max}$ and it is $O(n^0) = const$.
- c) the “worst” term of $(1 + \sum_m a_m q^{\sigma_m} n^{-\nu_{max}/|\mu_{max}| \cdot \sigma_m - \rho_m})^2$ is the one with $n^{-2\nu_{max}/|\mu_{max}| \cdot \sigma_{min} - \rho_{min}}$ (or the 1)

Denominator:

- a.i) $n^{-\nu_{max}/|\mu_{max}| \cdot 2\beta}$
- b.i) the dominant term of $(1 + \sum_m (max))^2$ is $n^{2(-\nu_{max}/|\mu_{max}| \cdot \sigma_{min} - \rho_{min})}$ (or the 1)
- a.ii) $n^{\nu_{min}/\mu_{min} \cdot (1-2\beta)}$
- b.ii) the dominant term of $(1 + \sum_m (min))^2$ is $n^{2(\nu_{min}/\mu_{min} \cdot \sigma_{max} - \rho_{max})}$ (or the 1)

If a.i) and b.i) dominate [i.e. $-\nu_{max}/|\mu_{max}| \cdot 2\beta + 2(-\nu_{max}/|\mu_{max}| \cdot \sigma_{min} - \rho_{min}) \geq \nu_{min}/\mu_{min} \cdot (1-2\beta) + 2(\nu_{min}/\mu_{min} \cdot \sigma_{max} - \rho_{max})$] then the expression will go to a constant, else it will go to 0.

2.term [$\int_c^\infty / (\int_0^c + \int_c^\infty)$] contains:

Numerator:

a) $n^{\nu_{min}/\mu_{min} \cdot (1-2\beta)}$

b) the “worst” term of $(\sum_k c_k \mu_k q^{\mu_k} n^{\nu_{min}/\mu_{min} \cdot \mu_k - \nu_k})^2$ is the one where $k = k_{min}$ and it is $O(n^0) = const.$

c) the “worst” term of $(1 + \sum_m a_m q^{\sigma_m} n^{\nu_{min}/\mu_{min} \cdot \sigma_m - \rho_m})^2$ is the one with $n^{2(\nu_{min}/\mu_{min} \cdot \sigma_{max} - \rho_{max})}$ (or the 1)

Denominator: same as above

If a.ii) and b.ii) dominate [i.e. $-\nu_{max}/|\mu_{max}| \cdot 2\beta + 2(-\nu_{max}/|\mu_{max}| \cdot \sigma_{min} - \rho_{min}) \leq \nu_{min}/\mu_{min} \cdot (1-2\beta) + 2(\nu_{min}/\mu_{min} \cdot \sigma_{max} - \rho_{max})$], then the expression will go to a constant, or else it will go to 0.

Since the two conditions for the two terms mutually exclude each other, except for equality, when both terms yield constants, there will always be one term going to zero and the other going to a constant. Therefore, the whole expression must contain at least one constant term.

To complete the proof one must show that this constant can not be cancelled exactly in the limit by negative constant terms. Unfortunately, there are 36 positive “pseudo-terms” (each containing many terms in the sums over k and m) and 30 negative ones. It would be very difficult to study these terms in detail to make sure that no total cancelation can occur. So, strictly speaking, the proof is not complete. From a practical point of view, however, there is enough evidence to safely assume this. Hence, $0 \notin spec(A)$ for $1/4 < \beta < 1/2$.

Case $0 < \beta \leq 1/4$

Start with an example and choose:

$$\psi_n := N_n(\frac{1}{2})^{-2\beta} (qs)^{-\beta} [(q^{-1} + s^{-1}) + i\beta^{-1}(p-r)]^{-2\beta} e^{-q/n-1/(qn)} e^{-(p-r)^2/n^2}$$

Without the regularization for small q (the $e^{-1/(qn)}$ factor), ψ_n would be square integrable. But, $A\psi_n = \{-q^2/n^2 + 2\beta[+]^{-1}1/n + (2-2\beta)q/n - \beta^2 q^{-2}(4ip/n^2[+]^{-1} + 4p^2/n^4 - 2/n^2) + \beta^2 q^{-1}8ip/n^2\}\psi_n$ would not be square integrable because of the terms with the factors q^{-2} and q^{-1} . One must have $\psi_n \in D(A)$, and this is why the regularization for small q was additionally introduced.

This complicates things, not only because there will be even more terms, but it will be hard to extract the correct n -dependence as well. (A regularization term like $e^{-1/(qn)}$ can not at the same time with $e^{-q/n}$ get rid of its n -dependence by a substitution like $q \rightarrow qn$, and that was the trick used above). The problem must be split in regions ($\int_0^c$ and $\int_c^\infty$).

$$\begin{aligned}
\|\psi_n\|^2 &\propto N_n^* N_n \int_0^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} \\
&\quad \times \int_{-\infty}^\infty dp [(q^{-1} + s^{-1})^2 + \beta^{-2} p^2]^{-2\beta} e^{-2p^2/n^2} \\
&= N_n^* N_n n^{1-4\beta} \int_0^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} \\
&\quad \times \int_{-\infty}^\infty dp [(q^{-1} + s^{-1})^2 n^{-2} + \beta^{-2} p^2]^{-2\beta} e^{-2p^2} \\
&= N_n^* N_n n^{1-4\beta} \left\{ \int_0^c dq e^{-2q/n-2/(qn)} q^{-2\beta} \times I_{(0,0)}(n) \right. \\
&\quad \left. + \int_c^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} \times I_{(0,0)}(n) \right\} \\
&\leq N_n^* N_n n^{1-4\beta} \left\{ n^{2\beta-1} \int_0^c dq e^{-2/q} q^{-2\beta} \cdot I_{(0,0)}^{small}(n, q) \right. \\
&\quad \left. + n^{1-2\beta} \int_c^\infty dq e^{-2q} q^{-2\beta} \cdot e^{-2/(cn^2)} I_{(0,0)}^{large}(n, q) \right\} \\
&\leq N_n^* N_n \{ n^{-2\beta} \cdot c_1 + n^{2-6\beta} \cdot c_2 \}
\end{aligned}$$

where (for reasons that become obvious later)

$$I_{(0,0)}(n, q) := \int_{-\infty}^\infty dp [(q^{-1} + s^{-1})^2 n^{-2} + \beta^{-2} p^2]^{-2\beta} e^{-2p^2}$$

$$I_{(0,0)}^{small}(n, q) := \int_{-\infty}^\infty dp [(nq^{-1} + s^{-1})^2 n^{-2} + \beta^{-2} p^2]^{-2\beta} e^{-2p^2} \xrightarrow{n \rightarrow \infty} I_{(0,0)}^{small}(q)$$

$$I_{(0,0)}^{large}(n, q) := \int_{-\infty}^\infty dp [((qn)^{-1} + s^{-1})^2 n^{-2} + \beta^{-2} p^2]^{-2\beta} e^{-2p^2} \xrightarrow{n \rightarrow \infty} I_{(0,0)}^{large} = const$$

$$c_1 := \int_0^c dq e^{-2/q} q^{-2\beta} \cdot I_{(0,0)}^{small}(q)$$

$$c_2 := \int_c^\infty dq e^{-2/q} q^{-2\beta} \cdot I^{large}$$

So $N_n^* N_n \geq \text{const} \cdot [c_1 n^{-2\beta} + c_2 n^{6\beta-2}]^{-1} \xrightarrow{n \rightarrow \infty} \text{const} \cdot n^{-2+6\beta}$.

Compute $A\psi_n$ and write “ p ” instead of “ $(p-r)$ ” since this will make no difference upon integration:

$$\begin{aligned} A\psi_n = & \{-q^2(1-q^{-2})n^{-2} + 2q[1-\beta(1-q^{-2})]n^{-1} \\ & - 4\beta^2 q^{-2} p^2 n^{-4} + 2\beta^2 q^{-2} n^{-2} + 4i\beta^2 q^{-1} p n^{-2} \\ & + 2\beta[+]^{-1}(1-q^{-2})n^{-1} + 4i\beta^2 q^{-2}[+]^{-1} p n^{-2}\} \psi_n \end{aligned}$$

The norm is:

$$\begin{aligned} \|A\psi_n\|^2 &= \int_0^\infty dq \int_{-\infty}^\infty dp (A\psi_n)^*(A\psi_n) \\ &= \int_0^\infty dq \int_{-\infty}^\infty dp \{[-q^2(1-q^{-2})n^{-2} + 2q[1-\beta(1-q^{-2})]n^{-1} \\ &\quad - 4\beta^2 q^{-2} p^2 n^{-4} + 2\beta^2 q^{-2} n^{-2}\}^2 + 16\beta^4 q^{-2} p^2 n^{-4} \\ &\quad + \{\dots\} 2\beta \frac{[+] + [-]}{[]} (1-q^{-2})n^{-1} + \{\dots\} 4\beta^2 q^{-2} p i \frac{[-] - [+]}{[]} n^{-2} \\ &\quad + 8\beta^3 q^{-1} (1-q^{-2}) p i \frac{[+] - [-]}{[]} n^{-3} + 16\beta^4 q^{-3} p^2 \frac{[+] + [-]}{[]} n^{-4} \\ &\quad + 4\beta^2 (1-q^{-2})^2 []^{-1} n^{-2} + 16\beta^4 q^{-4} p^2 []^{-1} n^{-4}\} \psi_n^* \psi_n \end{aligned}$$

where the definitions of $[+]$, $[-]$, $[]$ from above were used. $[+] + [-] = q^{-1} s^{-1}$, $\pm([-] - [+]) = \pm(2\beta^{-1} p)$. Here, $\{\dots\}$ are the “ q -terms” in the braces above.

Some of the terms will be explicitly calculated, the others will be discussed in words.

The first term is proportional to:

$$\begin{aligned}
& N_n^* N_n \int_0^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} (q^2 - 1)^2 n^{-4} \int_{-\infty}^\infty dp e^{-p^2/n^2} [\]^{-2\beta} \\
= & N_n^* N_n n^{1-4\beta} \int_0^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} (q^4 - 2q^2 + 1) n^{-4} I_{(0,0)}(n, q) \\
= & N_n^* N_n n^{1-4\beta} \left[\int_0^c dq e^{-2q/n-2/(qn)} q^{-2\beta} (q^4 - 2q^2 + 1) n^{-4} I_{(0,0)}(n, q) \right. \\
& \quad \left. + \int_c^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} (q^4 - 2q^2 + 1) n^{-4} I_{(0,0)}(n, q) \right] \\
\geq & N_n^* N_n n^{1-4\beta} \left[n^{2\beta-1} \int_0^c dq e^{-2c/n^2} e^{-2/q} q^{-2\beta} \right. \\
& \quad \times (n^{-4} q^4 - 2n^{-2} q^2 + 1) n^{-4} I_{(0,0)}^{small}(n, q) \\
& \quad \left. + n^{1-2\beta} \int_c^\infty dq e^{-2q} e^{-2/(cn^2)} q^{-2\beta} \right. \\
& \quad \left. \times (n^4 q^4 - 2n^2 q^2 + 1) n^{-4} I_{(0,0)}^{large}(n, q) \right]
\end{aligned}$$

Notice that the first part (small q) goes rapidly to zero. In the second part everything goes to zero, except the q^4 -term, which will go to a constant.

One can easily generalize this: Terms with a higher power of $1/n$ than of q will all go to zero (e.g. q/n^2 etc.) and will be called “more- n ” terms. Terms with the same power will go to a constant (e.g. q/n , q^2/n^2 etc.). Terms with $O(q^0) = O(1)$ will vanish in the limit since they come with some power of $1/n$. All these terms can be called “large q dominated” - since the positive powers of q help for small q .

Terms with a negative power of q (e.g. $1/(qn)$, $1/(qn^2)$, $1/(q^2 n^2)$ etc.) can be called “small q dominated”. It is for them that one needs the regularization for small q . (They help for large q , so the $\int_c^\infty$ -part will quickly go to zero). Here is an example:

$$\begin{aligned}
& N_n^* N_n \int_0^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} q^{-1} n^{-1} \int_{-\infty}^\infty dp e^{-p^2/n^2} []^{-2\beta} \\
&= N_n^* N_n n^{1-4\beta} \int_0^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} q^{-1} n^{-1} I_{(0,0)}(n, q) \\
&= N_n^* N_n n^{1-4\beta} \left[\int_0^c dq e^{-2q/n-2/(qn)} q^{-2\beta} q^{-1} n^{-1} I_{(0,0)}(n, q) \right. \\
&\quad \left. + \int_c^\infty dq e^{-2q/n-2/(qn)} q^{-2\beta} q^{-1} n^{-1} I_{(0,0)}(n, q) \right] \\
&\geq N_n^* N_n n^{1-4\beta} \left[n^{2\beta-1} \int_0^c dq e^{-2c/n^2} e^{-2/q} q^{-2\beta} q^{-1} I_{(0,0)}^{small}(n, q) \right. \\
&\quad \left. + n^{1-2\beta-2} \int_c^\infty dq e^{-2q} e^{-2/(cn^2)} q^{-2\beta} q^{-1} I_{(0,0)}^{large}(n, q) \right]
\end{aligned}$$

Because $N_n^* N_n \geq \text{const} \cdot [c_1 n^{-2\beta} + c_2 n^{6\beta-2}]^{-1}$ everything goes to zero. This can be generalized to all “small q dominated” terms.

The “ p ”- and “[]”-terms:

There are two kinds of terms involving p and the []-bracket. The first kind involves just []⁻¹, and so one sets:

$$\begin{aligned}
\int_{-\infty}^\infty dp e^{-p^2/n^2} []^{-2\beta} \cdot []^{-1} &= n^{1-4\beta} n^{-2} \int_{-\infty}^\infty dp e^{-p^2} []^{-2\beta-1} \\
&=: n^{1-4\beta} n^{-2} I_{(0,-2)}(n, q)
\end{aligned}$$

Now the meaning of the subscripts of the integrals I becomes clear. The 0 means no extra power of p , the -2 means one extra []⁻¹ (and the 2 rather than a 1 was chosen just because the []-bracket is quadratic in p and so one gets an extra n^{-2} from a simple []⁻¹). In the different regions, the $I_{(0,-2)}$ will go to $I_{(0,-2)}^{small}(n, q) \xrightarrow{n \rightarrow \infty} I_{(0,-2)}^{small}(q)$, and $I_{(0,-2)}^{large}(n, q) \xrightarrow{n \rightarrow \infty} I_{(0,-2)}^{large} = \text{const.}$

The second term involves p^2 and []⁻¹ and one sets:

$$\begin{aligned}
\int_{-\infty}^\infty dp e^{-p^2/n^2} p^2 []^{-2\beta} \cdot []^{-1} &= n^{1-4\beta} \int_{-\infty}^\infty dp e^{-p^2} p^2 []^{-2\beta-1} \\
&=: n^{1-4\beta} I_{(2,-2)}(n, q)
\end{aligned}$$

Compared to the original term without the extra factors, there is no different n -dependence, only the integral changed [$I_{(2,-2)}(n, q)$ instead of $I_{(0,0)}(n, q)$].

So all terms involving the extra p - and $[\]$ -factors are of the “small q dominated” or “more- n ” type and will vanish in the limit. So the only terms going to constants are the “large q dominated”, non-“more- n ” terms which already appeared in the easier case $1 < 4\beta < 2$ as well.

The conclusion is again that the norm of $A\psi_n$ will not vanish for $n \rightarrow \infty$ (the same comment concerning cancelations as in the case $1/4 < \beta < 1/2$ applies).

Terms coming from a p -regularization used to vanish in the limit, so one does not expect new things to happen for a general ψ_n , and the previous discussion of the general case (for $1/4 < \beta < 1/2$) is enough to convince for $0 < 4\beta \leq 1$ as well.

Note that the case $4\beta = 1$ as a limiting case is well behaved, namely the norm-squares of ψ_n go to $n^{-1/2}$ {for $4\beta \searrow 1$: $N_n^* N_n \propto n^{2\beta-1}$ and for $4\beta \nearrow 1$: $N_n^* N_n \geq \text{const} \cdot [c_1 n^{-2\beta} + c_2 n^{6\beta-2}]^{-1} \xrightarrow{n \rightarrow \infty} \text{const} \cdot n^{-2+6\beta}$ }

A.4 Alternative computation of the denominator of Eq. (75)

Another way to find the ν -dependence of the denominator of Eq. (75) is the following. Use the definition of the Whittaker function [Eq. (67)] $W_{\mu,\nu}(z) = e^{-z/2} z^{1/2+\nu} U(1/2 - \mu + \nu, 1 + 2\nu, z)$ and of the confluent hypergeometric function $U(a, b, z) = \Gamma^{-1}(a) \int_0^\infty e^{-zt} t^{a-1} (1+t)^{b-a-1} dt$ applied to $\mu = 1/2$, $\nu = i\lambda$ and extract the the different orders of λ directly. Attention is required because the integral definition demands a real part for a , which is introduced as ϵ , and all expressions are computed with the complex variable $\xi := \epsilon + i\lambda$. The limit $\epsilon \rightarrow 0$ is taken in the end (or when not critical). Although this approach to the problem is different from the one before, it does not come as a surprise that there is a need for an auxiliary parameter ϵ . In the first approach, the critical point was the evaluation of the hypergeometric series after the x -integration. Here, it is the definition of the confluent hypergeometric function U before the x -integration.

The wave function becomes:

$$\begin{aligned} x^{1/2} \psi'_\lambda(x) &= x^{1/2} x^{-1/2} \psi_\lambda(\ln(x)) \\ &= \left(\frac{\lambda \sinh(2\pi\lambda)}{\pi^2} \right)^{1/2} \Gamma(i\lambda) x^{-1/2} e^{-x/2} x^{1/2+i\lambda} \Gamma^{-1}(i\lambda) \int_0^\infty e^{-xt} t^{-1+\xi} (1+t)^\xi dt \end{aligned}$$

Notice that the small real part has been added only in the integrand, leading to a somewhat mixed notation.

This comes together with its complex conjugate. So, by omitting the pure λ -dependent factors for a while, one gets an integral

$$\begin{aligned}
& \lim_{\epsilon \rightarrow 0} \int_0^\infty dx e^{-x} \left\{ \int_0^\infty e^{-xt} t^{-1+\xi} (1+t)^\xi dt \right\} \left\{ \int_0^\infty e^{-xt} t^{-1+\xi^*} (1+t)^{\xi^*} dt \right\} \\
&= \lim_{\epsilon \rightarrow 0} \int_0^\infty dx e^{-x} \left\{ \int_0^\infty e^{-xt} t^{-1+\xi} \left[1 + \ln(1+t)\xi + \frac{1}{2} \ln^2(1+t)\xi^2 + \dots \right] dt \right\} \\
&\quad \times \left\{ \int_0^\infty e^{-xt} t^{-1+\xi^*} \left[1 + \ln(1+t)\xi^* + \frac{1}{2} \ln^2(1+t)\xi^{*2} + \dots \right] dt \right\}
\end{aligned}$$

The next step is to interchange the sum, from the power series expansion, and the integral. This is only justified if one can make sure the existence of all integrals (to all orders) and the existence of the limit $\epsilon \rightarrow 0$. The latter is trivial, except for the summands with a “1”. Higher orders come at least with one factor of $\ln(1+t)$, which goes linearly to 0 for $t \rightarrow 0$, thus cancelling the otherwise critical t^{-1} .

The lowest order (take the 1’s from both braces) is:

$$\begin{aligned}
& \lim_{\epsilon \rightarrow 0} \int_0^\infty dx e^{-x} \left\{ \int_0^\infty e^{-xt} t^{-1+\xi} dt \right\} \left\{ \int_0^\infty e^{-xt} t^{-1+\xi^*} dt \right\} \\
&= \lim_{\epsilon \rightarrow 0} \int_0^\infty dx e^{-x} x^{-2\epsilon} \left\{ \int_0^\infty e^{-t} t^{-1+\xi} dt \right\} \left\{ \int_0^\infty e^{-t} t^{-1+\xi^*} dt \right\} \\
&= \lim_{\epsilon \rightarrow 0} \Gamma(1-2\epsilon) \Gamma(\xi) \Gamma(\xi^*) = \Gamma(i\lambda) \Gamma(-i\lambda) = O(1/\lambda^2)
\end{aligned}$$

Notice that the terminology “lowest order” refers to the expansion of $(1+t)^\xi$ only. Naturally, this “lowest order” term contains all possible powers of λ , but is the only term to really contain the lowest order which is $1/\lambda^2$. For the “higher orders”, the non-critical limit can be taken first and the expression is now

$$\begin{aligned}
& \int_0^\infty dx e^{-x} \left[\left\{ \int_0^\infty e^{-xt} t^{-1+i\lambda} [\mathbf{1} + \ln(1+t)i\lambda + \frac{1}{2} \ln^2(1+t)(i\lambda)^2 + \dots] dt \right\} \right. \\
& \quad \times \left\{ \int_0^\infty e^{-xt} t^{-1-i\lambda} [\mathbf{1} + \ln(1+t)(-i\lambda) + \frac{1}{2} \ln^2(1+t)(-i\lambda)^2 + \dots] dt \right\} \\
& \quad \left. - \text{“}\mathbf{1}, \mathbf{1} - \text{term”} \right] \\
& = \int_0^\infty dx e^{-x} \left[\left\{ \int_0^\infty e^{-xt} t^{-1} [1 + i \ln(t)\lambda + \frac{1}{2} (i \ln(t))^2 \lambda^2 + \dots] \right. \right. \\
& \quad \times [\mathbf{1} + \ln(1+t)i\lambda + \frac{1}{2} \ln^2(1+t)(i\lambda)^2 + \dots] dt \left. \right\} \\
& \quad \times \left\{ \int_0^\infty e^{-xt} t^{-1} [1 - i \ln(t)\lambda + \frac{1}{2} (-i \ln(t))^2 \lambda^2 + \dots] \right. \\
& \quad \times [\mathbf{1} + \ln(1+t)(-i\lambda) + \frac{1}{2} \ln^2(1+t)(-i\lambda)^2 + \dots] dt \left. \right\} \\
& \quad \left. - \text{“}\mathbf{1}, \mathbf{1} - \text{term”} \right]
\end{aligned}$$

where $-\text{“}\mathbf{1}, \mathbf{1} - \text{term”}$ indicates the term already taken care of. This is just a notation to avoid having to write out everything. The only thing to prove is the existence of all integrals. They are of the type

$$\begin{aligned}
& \int_0^\infty dx e^{-x} \left[\left\{ \int_0^\infty e^{-xt} t^{-1} [\ln(t)]^n [\ln(1+t)]^m dt \right\} \right. \\
& \quad \times \left. \left\{ \int_0^\infty e^{-xs} [\ln(s)]^{\tilde{n}} [\ln(1+s)]^{\tilde{m}} ds \right\} \right] \quad (108)
\end{aligned}$$

where $m, \tilde{m} \in \mathbb{N}$, $n, \tilde{n} \in \mathbb{N}_0$.

Look at the integral $\int_0^\infty e^{-xt} [\ln(t)]^n [\ln(1+t)]^m dt$ first. Rewrite it as $\int_0^\infty = \int_0^c + \int_c^\infty$ where $c > e - 1$; e is the Euler number.

$$\int_0^c e^{-xt} t^{-1} [\ln(1+t)]^m [\ln(t)]^n dt = f(x) \quad (109)$$

The “worst case” is $m = 1$, but $[\ln(t)]^n$ is integrable at 0 since $\ln(1+t) \approx t$ for small t , which takes care of the t^{-1} . Therefore, there is no problem for any x . Now, $f(x)$ is a constant for $x = 0$ and goes to 0 for $x \rightarrow \infty$. Since it is continuous, it is thus bounded, and so $f(x) \leq c_1 = \text{const}$.

$$\int_c^\infty e^{-xt} t^{-1} [\ln(1+t)]^m [\ln(t)]^n dt = g(x) \quad (110)$$

Since $c > e - 1 > 1$ the function $g(x)$ is certainly positive and $g(x)$ goes to 0 for large x . For $0 < \delta < 1$, the value of the function $g(x = \delta)$ is finite. Because $g(x)$ is continuous, it is bounded on the interval $[\delta, \infty)$ with bound d_1 . The integral does not exist for $x = 0$, so one has to extract the x -dependence for small x to determine x -integrability at 0 (in the following assume at least $x < 1$ so that $-\ln(x) > 0$). Since $c > e - 1$, there is a $k \in \mathbb{N}$ such that $[\ln(1+t)]^m \leq [\ln(t)]^{m+k}$. So Eq. (110) becomes

$$\begin{aligned}
& \int_c^\infty e^{-xt} t^{-1} [\ln(1+t)]^m [\ln(t)]^n dt \\
& \leq \int_c^\infty e^{-xt} t^{-1} [\ln(t)]^{n+m+k} dt = \int_{cx}^\infty e^{-t} t^{-1} [\ln(t/x)]^{n+m+k} dt \\
& = \int_{cx}^\infty e^{-t} t^{-1} [-\ln(x) + \ln(t)]^{n+m+k} dt \\
& = \int_{cx}^\infty e^{-t} t^{-1} \left\{ \sum_{l=0}^{n+m+k} \binom{n+m+k}{l} [-\ln(x)]^l [\ln(t)]^{n+m+k-l} \right\} dt \\
& \leq \int_{cx}^\infty e^{-t} t^{-1} \left\{ \sum_{l=0}^{n+m+k} \binom{n+m+k}{l} [-\ln(x)]^l |\ln(t)|^{n+m+k-l} \right\} dt \\
& \leq \sum_{l=0}^{n+m+k} \binom{n+m+k}{l} [-\ln(x)]^l \int_x^\infty e^{-t} t^{-1} |\ln(t)|^{n+m+k-l} dt \\
& =: \sum_{l=0}^{n+m+k} \binom{n+m+k}{l} [-\ln(x)]^l \cdot h_l(x) \tag{111}
\end{aligned}$$

Examine h_l alone for a moment:

$$\begin{aligned}
h_l(x) &= \int_x^\infty e^{-t} t^{-1} |\ln(t)|^{n+m+k-l} dt \\
&= \int_x^1 (\dots) + \int_1^\infty (\dots) \\
&\leq \int_x^1 t^{-1} |\ln(t)|^{n+m+k-l} dt + \text{const} \\
&= \int_{\ln(x)}^0 |u|^{n+m+k-l} du + \text{const} = \int_{\ln(x)}^0 (-u)^{n+m+k-l} du + \text{const} \\
&= (n+m+k-l+1)^{-1} (-1)^{n+m+k-l+1} [\ln(x)]^{n+m+k-l+1} + \text{const} \\
&= (n+m+k-l+1)^{-1} [-\ln(x)]^{n+m+k-l+1} + \text{const}
\end{aligned}$$

Putting this in the last line of Eq. (111), one sees that, for small x , the function $g(x)$ is dominated by a polynomial in $-\ln(x)$ of degree $(n + m + k + 1)$. This is named $\text{Poly}^{n+m+k+1}[-\ln(x)]$.

The same arguments hold for the s -integral in Eq. (108). Denote the new constants in an obvious way by c_2 , d_2 and $\tilde{k}$ and get functions $\tilde{f}$, $\tilde{g}$, $\tilde{h}_l$ at an intermediate step, and $\text{Poly}^{\tilde{n}+\tilde{m}+\tilde{k}+1}[-\ln(x)]$. The integral (108) is bounded by

$$\begin{aligned}
& \int_0^\infty e^{-x}[f(x) + g(x)][\tilde{f}(x) + \tilde{g}(x)]dx \\
& \leq \int_0^\infty e^{-x}[c_1 + g(x)][c_2 + \tilde{g}(x)]dx \\
& = \int_0^\epsilon e^{-x}[c_1 + g(x)][c_2 + \tilde{g}(x)]dx + \int_\epsilon^\infty e^{-x}[c_1 + g(x)][c_2 + \tilde{g}(x)]dx \\
& \leq \int_0^\epsilon e^{-x}\{c_1 + \text{Poly}^{n+m+k+1}[-\ln(x)]\}\{c_2 + \text{Poly}^{\tilde{n}+\tilde{m}+\tilde{k}+1}[-\ln(x)]\}dx \\
& \quad + \int_\epsilon^\infty e^{-x}[c_1 + d_1][c_2 + d_2]dx
\end{aligned}$$

This expression is smaller than infinity since $[\ln(x)]^r$ is integrable at 0 for any r , and e^{-x} takes care of the integrability at infinity.

One could go on and calculate the numerical factors for the different orders of λ , i.e. develop the prefactor $\lambda \sinh(2\pi\lambda)/\pi^2$ and the Gamma functions $\Gamma(i\lambda)\Gamma(-i\lambda)$ in power series, compute the relevant integrals for each power, and collect the numerical coefficients belonging to that power. Surprisingly, the coefficients are not the same as in Eq. (76). But, they do not have to be, since the two approaches to the problem used different regularizations to deal with only conditionally convergent expressions. The final result does not depend on these coefficients since a quotient is taken.

B Acknowledgments

Thanks are expressed to Prof. Dr. A. Schenzle as first referee of this Diplomthesis and for his support of it, to Prof. Dr. J. von Delft for taking on the task of second referee, to the Deutsche Studienstiftung for financial support and encouragement, to the University of Florida and the members of its physics department for hospitality and to S. Shabanov and W. Bomstad for comments and helpful remarks.

Very special thanks go to Prof. Dr. J.R. Klauder who suggested the topic of this article - weak coherent states - which is based on his outstanding lifelong work. The research for this thesis was conducted under his supervision and could not have been done without his being a constant source of information as well as of ideas and encouragement at all times. This made working with him not only an honor but a great pleasure, too.

References

- [1] P.A.M. Dirac, *The Principles of Quantum Mechanics*, Oxford University Press, Oxford, Third Edition, 1947, p. 114
- [2] R. Shankar: *Principles of Quantum Mechanics*, Plenum, New York, Exercise 7.4.10, pages 225-227
- [3] E. Wigner, *On the Quantum Correction For Thermodynamic Equilibrium*, Phys. Rev. 40, 749-759, 1932
- [4] E. Wigner, *Quantum-Mechanical Distribution Functions Revisited*
- [5] J.E. Moyal, *Quantum Mechanics as a Statistical Theory*, Proc. Camb. Phil. Soc. 45, 99-124, 1949
- [6] G.E. Baker, *Formulation of Quantum Mechanics Based on the Quasi-Probability Distribution Induced on Phase Space*, Phys. Rev. 109, No. 6, 2198-2206, 1958
- [7] D.B. Fairlie, *The formulation of quantum mechanics in terms of phase space functions*, Proc. Comb. Phil. Soc. 60, 581-586, 1963
- [8] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, *Quantum Mechanics as a Deformation of Classical Mechanics*, Lett. Math. Phys. 1, 521-530, 1977
- [9] T. Curtright, D. Fairlie, C. Zachos, *Features of the time-independent Wigner functions*, Phys. Rev. D 58, 1998
- [10] T. Curtright, T. Uematsu, C. Zachos, *Generating all Wigner functions*, J. Math. Phys. 42, No. 6, 2396-2415, 2001
- [11] C. Zachos, *Deformation Quantization: Quantum Mechanics Lives and Works in Phase Space*, International Journal of Modern Physics A, Vol. 17, No. 3, 297-316, 2002
- [12] J. Śniatycki, *Geometric Quantization and Quantum Mechanics*, Springer, New York, 1980
- [13] J.R. Klauder, E.C.G. Sudarshan, *Fundamentals of Quantum Optics*, W.A. Benjamin, New York, 1968
- [14] J.R. Klauder, B-S. Skagerstam, *Coherent States, Applications in Physics and Mathematical Physics*, World Scientific, Singapore, 1985

- [15] A.M. Perelomov, *Coherent States for Arbitrary Lie Group*, Commun. math. Phys. 26, 222-236, 1972
- [16] J.R. Klauder, *Continuous-Representation Theory. I. Postulates of Continuous-Representation Theory*, J. Math. Phys. 4, No. 8, 1055-1058, 1963
- [17] J.R. Klauder, *Continuous Representation Theory. II. Generalized Relation between Quantum and Classical Dynamics*, J. Math. Phys. 4, No. 8, 1058-1073, 1963
- [18] J.R. Klauder, *Continuous Representation Theory. III. On Functional Quantization of Classical Systems*, J. Math. Phys. 5, 177-187, 1964
- [19] R.H. Cameron, *A Family of Integrals Serving to Connect the Wiener and Feynman Integrals*, Journal of Math. and Phys. 39, 126-140, 1960
- [20] R.H. Cameron, *The Ilstow and Feynman Integrals*, J. Analyse Math. 10, 287-361, 1962/63
- [21] P. Chernoff, *Note on Product Formulas for Operator Semigroups*, J. Func. Analysis 2, 238-242, 1968
- [22] J.R. Klauder, *Continuous Representations and Path Integrals, Revisited*, reprinted in *Path Integrals* edited by G.J. Papadopoulos and J.T. Devreese, Plenum Publishing Corporation, 1978
- [23] H. Meschkowski, *Hilbertsche Räume mit Kernfunktion*, Springer, 1962
- [24] E.C.G. Sudarshan, *Equivalence of Semiclassical and Quantum Mechanical Descriptions of Statistical Light Beams*, Phys. Rev. Lett. 10, 277-279, 1963
- [25] E.W. Aslaksen, J.R. Klauder, *Continuous Representation Theory Using the Affine Group*, J. Math. Phys. 10, 2267-2275, 1969
- [26] I. Daubechies, J.R. Klauder, T. Paul, *Wiener measures for path integrals with affine kinematic variables*, J. Math. Phys 28, 85-102, 1986
- [27] J.R. Klauder, *Coherent State Path Integral without Resolutions of Unity*, Foundations of Physics 31, No. 1, 57-67, 2001
- [28] J.R. Klauder, *Quantization Is Geometry, after All*, Annals of Physics 188, 120-141, 1988

- [29] G. Roepstorff, *Path Integral Approach to Quantum Physics, An Introduction*, Springer, 1994
- [30] C. Grosche, F. Steiner, *Handbook of Feynman Path Integrals*, Springer Tracts in Modern Physics 145, Springer, 1998
- [31] H. Goldstein, *Classical Mechanics*, Addison-Wesley, Cambridge, MA, 1950
- [32] R. Abraham, J. Marsden, *Foundations of Mechanics*, Benjamin, New York, 1978
- [33] J.R. Klauder, *Understanding Quantization*, Foundations of Physics 27, No. 11, 1467-1483, 1997
- [34] B. Simon, *Functional Integration and Quantum Mechanics*, Academic Press, 1979
- [35] A.V. Skorokhod, *Studies in the Theory of Random Processes*, Addison Wesley, 1965
- [36] R.P. Feynman, *The Principle of Least Action in Quantum Mechanics*, Ph. D. thesis, Princeton University, 1942
- [37] J.R. Klauder, *Quantization Is Geometry, after All*, Annals of Physics 188, 120-141, 1988
- [38] J.R. Klauder, *Path integrals and stationary-phase approximations*, Phys. Rev. D, 19, No.8, 2349-2356, 1979
- [39] I. Daubechies, J.R. Klauder, *Quantum-mechanical path integrals with Wiener measure for all polynomial Hamiltonians. II*, J. Math. Phys. 26, No.9, 2239-2256, 1985
- [40] P.M. Morse, *Diatomic Molecules According to the Wave Mechanics. II. Vibrational Levels*, Phys. Rev. 34, 57-64, 1929
- [41] P. Maraner, *Landau Ground State on Riemannian Surface*, Modern Physics Letters A, Vol. 7, No. 27, 2555-2558, 1992
- [42] J.R. Klauder, *Noncanonical quantization of gravity. I. Foundations of affine quantum gravity*, J. Math. Phys. 40, No. 11, 1999
- [43] J.R. Klauder, *Noncanonical quantization of gravity. II. Constraints and the physical Hilbert space*, J. Math. Phys. 42, No. 9, 2001

- [44] D.W. Stroock, *Dirichlet forms*, Bulletin of the American Mathematical Society 33, No. 1, 87-92, 1996
- [45] I.S. Gradshteyn, I.M. Ryzhik, *Tables of integrals, series and products*, Academic Press, San Diego, 1980
- [46] C. Grosche, *The Path Integral on the Poincaré Upper Half-Plane with a Magnetic Field and for the Morse Potential*, Annals of Physics 187, 110-134, 1988
- [47] M. Abramowitz, *Handbook of Mathematical Functions*, Dover, 1965
- [48] J. Blank, P. Exner, M. Havlíček, *Hilbert Space Operators in Quantum Physics*, AIP Series in Computational and Applied Mathematical Physics, New York, 1994
- [49] S. Shabanov, personal communications; work unpublished